

## INVITED ARTICLE

# Adolescent Idiopathic Scoliosis

Jayesh Trivedi

**How to cite this article:** Trivedi J. Adolescent Idiopathic Scoliosis. *J Postgrad Med Edu Res* 2017;51(2):58-67.

**Source of support:** Nil

**Conflict of interest:** None

## INTRODUCTION

Traditionally idiopathic scoliosis is divided into three categories according to the age of onset: Infantile (0–3 years), juvenile (4–9 years), and adolescent (10 years to skeletal maturity). Adolescent idiopathic scoliosis (AIS) is a structural lateral curvature of the spine measuring at least 10° of Cobb's angle and occurring at or near the onset of puberty. The prevalence of adolescent scoliosis in the adolescent population is 2 to 3%.<sup>1-4</sup> However, as the curve magnitude increases, the prevalence decreases. The larger curves have a female preponderance in a ratio of 4:1. In the smaller curves, males and females have almost an equal prevalence.<sup>3,4</sup> The prevalence of curves greater than 30° is about 0.2%, and it is less than 0.1% for curves greater than 40°.<sup>3,4</sup>

## ETIOLOGY

As the term idiopathic indicates, the etiology of adolescent scoliosis is unknown. However, hereditary, environmental, collagen and connective tissue defects, neuromuscular and biomechanical factors have all been indicated in the etiology of adolescent scoliosis.

### Hereditary Factors

It has long been known that hereditary factors play a role in the etiology of idiopathic scoliosis (IS).<sup>5</sup> Inheritance of scoliosis in five generations was described by Garland.<sup>6</sup> In a cohort of 1,463 individuals with IS, Grauers et al found that among those treated with a brace or surgery

for scoliosis, 53% reported one or more relatives with scoliosis, compared to 46% of the untreated, pointing towards a slightly higher risk of treatment in the presence of a family history of scoliosis.<sup>7</sup>

### Endocrine Factors

Thillard discovered that pinealectomized chickens developed scoliosis.<sup>8,9</sup> This was repeated in bipedalised rats and a deficiency of melatonin was suggested to be causative of IS.<sup>10,11</sup> Further studies, however, showed that adolescent IS patients had normal melatonin levels and pinealectomized monkeys did not develop scoliosis.<sup>12</sup>

Calmodulin is a calcium-binding receptor protein which regulates contractile properties in platelets and muscles, and also interacts with melatonin. Increased levels of calmodulin in platelets and an asymmetrical distribution of calmodulin in paraspinal muscles compared to healthy controls have been described in IS patients.<sup>13</sup>

### Biomechanical Factors

Dickson et al showed that vertebral bodies were wedged in the sagittal plane in IS patients, causing an apical lordosis in thoracic curvatures. They suggested that lordosis, in a region, i.e., normally kyphotic, creates a rotational moment in the spine thus causing scoliosis.<sup>14</sup>

### Neuromuscular Factors

Neuromuscular conditions producing an asymmetry of the transverse-spinalis muscle, abnormality in visual, vestibular, proprioceptive and postural control have also been implicated in AIS.<sup>15</sup> In addition, tonsillar ectopia with abnormal somato-sensory evoked potentials, larger foramen magnum, and left-right brain asymmetries point to neural origin of AIS.<sup>16,17</sup> These findings explain a poor performance in combined visual and proprioception, as well as spatial orientation tests and impaired postural balance in AIS patients.

The higher prevalence of AIS observed in females may be due to the fact that girls attain adolescent skeletal growth spurt in relative postural immaturity, compared to boys who go through their pubertal rapid growth at later age when their posture is more mature. Spine slenderness and ectomorphy are other risk factors in girls.<sup>18</sup>

Consultant Spine Surgeon

Centre for Spinal Studies, Robert Jones and Agnes Hunt Orthopedics Hospital, Oswestry, Shropshire, United Kingdom

**Corresponding Author:** Jayesh Trivedi, Consultant Spine Surgeon, Centre for Spinal Studies, Robert Jones and Agnes Hunt Orthopedics Hospital, Oswestry, Shropshire, United Kingdom, Phone: +441691404663, e-mail: jayeshmtrivedi@aol.com

## NATURAL HISTORY OF ADOLESCENT IDIOPATHIC SCOLIOSIS

The main concerns are that patients with IS relate mainly to curve progression, occurrence of low back pain, and pulmonary function.

### Curve Progression

In the immature patient, factors affecting curve progression relate mainly to curve factors and growth potential. The majority of information on curve progression relates mainly to scoliosis in girls. Six factors influencing curve progression have been identified. These include:

1. The younger the patient at the time of diagnosis, the more risk of progression.<sup>4,19</sup>
2. The greatest risk of progression happens just prior to the onset of menarche.
3. Risser grades 0 and 1 are more likely to progress than the higher Risser grades.<sup>19</sup>

Joseph C Risser first described what now is called the Risser sign in 1958. Risser observed that the state of

ossification of the iliac apophysis was associated with the state of a patient's spinal skeletal maturity.<sup>20</sup>

In the US system, the ossification of the iliac apophysis is divided into quarters, whereas in the European system the iliac apophysis ossification is divided into thirds with the fourth stage representing the commencement of fusion of the apophysis to the crest posteriorly<sup>21</sup> (Fig. 1).

Lonstein et al collated the risk of progression based on Risser grade and curve magnitude<sup>19</sup> (Table 1).

4. Males with comparable curves have one-tenth the risk of progression compared to females.<sup>4,22</sup>
5. Double curves have a greater chance of progression. Lumbar curves more than 30°, and where the fifth lumbar vertebra is high riding are more likely to progress.<sup>23</sup>
6. Finally, curves with higher magnitude are more likely to progress after skeletal maturity. Curves between 50 to 75° are likely to progress at 0.5 to 1.0° per year.<sup>23</sup>

Sanders et al modified the Tanner-Whitehouse staging system to assess risk of progression in adolescent scoliosis. They emphasized the use of the physes




Fig. 1: Depiction of Risser sign

**Table 1:** Risk of progression

Risser	Curve magnitude 5–19	Curve magnitude 20–25
0–1	22%	68%
2–4	1.6%	23%

**Table 2:** Skeletal maturity based on hand physes

Stage	Key features
1 Juvenile slow	Digital epiphyses are not covered.
2 Preadolescent slow	All digital epiphyses are covered.
3 Adolescent rapid—early	The preponderance of digits are capped. The second through fifth metacarpal epiphyses are wider than their metaphyses.
4 Adolescent rapid—late	Any of distal phalangeal physes are clearly beginning to close.
5 Adolescent steady—early	All distal phalangeal physes are closed and others are open.
6 Adolescent steady—late	Middle or proximal phalangeal physes are closing.
7 Early mature	Only distal radial physis is open. Metacarpal physes scars may be present.
8 Mature	Distal radial physis is completely closed.

of the hand to grade the skeletal maturity into eight stages (Table 2).<sup>24</sup>

## Back Pain

The incidence of back pain in the general population is reported at between 60 and 80%.<sup>25</sup> Some studies have indicated that the prevalence of back pain in adolescents with IS are similar to the general adolescent population while other studies concluded that adolescents with IS experience more back pain and more severe back pain than their peers.<sup>26,27</sup> Ramirez et al<sup>28</sup> reported a retrospective study of 2,442 patients (6–20 years) with IS showing that the lifetime prevalence of back pain was around 23% and the point prevalence of back pain was 9%. He concluded that these results were similar to those reported by other authors in the general pediatric and adolescent population. Dickson et al also reported that back pain was significantly more prevalent among the 165-subjects with scoliosis compared to the 100 age- and sex-matched control subjects (73 vs 52%) using a questionnaire survey (average age at survey was 17 years). Sato et al,<sup>29</sup> in an epidemiological study of over 43,000 adolescent school children in Japan the point and lifetime prevalence in Scoliosis group were 27.5 and 58.8% respectively, which both were significantly higher than in no scoliosis group. The pupils in scoliosis group also experienced significantly more back pain, a longer duration of back pain, and more recurrences of back pain in comparison to those without scoliosis.

## Pulmonary Function

Findings in pulmonary function studies in patients with untreated AIS demonstrate that only in patients with thoracic curves is there a direct correlation between decreased pulmonary function and increasing curve severity. Vital capacity and forced expiratory volume in 1 second decreases as the severity of thoracic curves increases.<sup>30–33</sup> The pattern of pulmonary disease in affected patients is, uniformly, restrictive lung disease. Smokers are generally affected much more severely than nonsmokers. Results in most studies show that significant limitations of forced vital capacity in nonsmokers do not occur until the curve approaches 100 to 120°. <sup>34</sup> However, the presence of thoracic hypo-kyphosis increases the loss of pulmonary function associated with curve severity.

Kim et al,<sup>35</sup> in a study of pulmonary function in adolescent scoliosis relative to the surgical procedure demonstrated a significant negative correlation between the preoperative Cobb's angle and percent predicted pulmonary function test values and a significant negative correlation between the number of involved vertebrae in the major curve and percent-predicted pulmonary function test values. The smaller number of involved vertebrae in the major curve had a strong correlation with higher pulmonary function test values preoperatively and hence, a better pulmonary function in Lenke 5 curves.

## CLASSIFICATION OF ADOLESCENT SCOLIOSIS

In 1983, Howard King presented his classification system for AIS. This system was based on the experience of John Moe in the surgical treatment of AIS patients with Harrington rod instrumentation. Curves were divided into five types and guidelines and recommendations for which levels should be instrumented were given according to those different curve types to preserve motion as much as possible.<sup>36</sup>

King and Moe defined five curve types:

*Type I:* An "S" shape deformity, in which both curves are structural and cross the CSVL, with the lumbar curve being larger than the thoracic one.

*Type II:* An "S" shape deformity, in which both curves are structural and cross the CSVL, with the thoracic curve being larger or equal to the lumbar one.

*Type III:* Major thoracic curve in which only the thoracic curve is structural and crosses the CSVL.

*Type IV:* Long "C" shape thoracic curve in which the fifth lumbar vertebra is centered over the sacrum and the fourth lumbar vertebra is tilted into the thoracic curve.

*Type V:* Double thoracic curve.

The drawbacks of the King classification was the failure to address the sagittal profile of the deformity and the low inter and intra-observer reliability.<sup>37,38</sup>

In an attempt to address these drawbacks, Lenke et al<sup>37</sup> proposed a classification, which is now widely used. This classification uses the following terminologies:

- Major curve: The curve of greatest magnitude and is always structural.
- Minor curve: A smaller curve which may be structural or nonstructural.
  - Nonstructural curve: A curve which bends to less than 25° on side bending radiographs.

According to these definitions, there are six different curve types (Fig. 2).

*Type I:* Main thoracic (MT) is the only structural curve while the others (proximal thoracic and lumbar or thoracolumbar) are nonstructural.

*Type II:* Double thoracic in which the MT is the major curve, the proximal thoracic (PT) is the minor curve but is structural, and the thoracolumbar (TL) or lumbar curves are minor and nonstructural.

*Type III:* Double major curve pattern in which the MT is the major curve and the lumbar is the minor one but is structural, whereas the PT is nonstructural.

*Type IV:* Triple major curve pattern when the MT is the major curve but all three curves are structural.

*Type V:* The TL or lumbar curve is the major and only structural curve, with the PT and/or MT curves being minor and nonstructural.

*Type VI:* The TL or lumbar curve is the major curve measuring at least 5° more than the MT which is the minor but structural curve.

To these basic six curve types, the lumbar spine modifier is added. This modifier is defined by the location of

Curve type (1-6)						
Lumbar spine modifier	Type 1 (main Thoracic)	Type 2 (double Thoracic)	Type 3 (double major)	Type 4 (triple major)	Type 5 (TL/L)	Type 6 (TL/L MT)
(No to minimal curve)	 1A*	 2A*	 3A*	 4A*		
(Moderate Curve)	 1B*	 2B*	 3B*	 4B*		
(Large curve)	 1C*	 2C*	 3C*	 4C*	 5C*	 6C*
Possible sagittal structural criteria (to determine Specific curve type)	Normal	PT kyphosis	TL syphosis	PT+TL kyphosis		

Fig. 2: Pictorial depiction of Lenke classification<sup>37</sup>




Fig. 3: Lumbar modifiers in Lenke classification

the CVSL on the apical vertebra of the lumbar curve. Lenke defined three lumbar modifiers (Fig. 3).

In addition, weightage is given to the sagittal profile of the curve by introducing the concept of a thoracic kyphosis modifier as follows:

- + (plus): When thoracic kyphosis measures  $>40^\circ$ .
- N (normal): When thoracic kyphosis measures between  $10$  and  $40^\circ$ .
- - (minus): When thoracic kyphosis measures  $<10^\circ$ .

### ASSESSMENT OF SCOLIOSIS

A detailed history is essential with respect to age of onset of deformity, menarche, and family history as well as the presence and absence of back pain. In addition the history should cover symptoms related to other systems including the central nervous system (developmental milestones, visual/hearing problems, gait abnormalities), cardiovascular system (cardiac murmurs), and the genitourinary system.

The physical examination should include an observation of presence or absence of secondary sexual characteristics to determine the status of skeletal maturity, and therefore, the risk of progression of the deformity. Gait should be examined for any evidence of neurological deficit or limb length discrepancy. The examination of the spine should include any skin discoloration (café-au-lait spots: Fig. 4), shoulder levels, contour of the spine, scapular, rib prominence, and symmetry of the waist creases (Figs 5A and B). A plumb line observation is done to see whether the head is aligned over the sacrum (a balanced spine, Fig. 6). The Adam's forward bend test is performed to assess for the rib prominence (Fig. 5B) and side bending in this position is used to assess for the flexibility of the curve. Alternately, the patient is placed prone and flexibility of the curve assessed manually by pushing against the rib prominence. A neurological examination is then undertaken.

Erect radiographs of the spine in PA and lateral projections are undertaken to measure the magnitude of the curve using the Cobb's angle and determine the Risser status and the status of the triradiate cartilages on the radiographs. Side bending PA radiographs of the spine





Fig. 4: Arrow indicates café-au-lait spot

will give an indication of the flexibility of the curve and help determine the extent of fusion at surgery.


### NONOPERATIVE TREATMENT OF ADOLESCENT SCOLIOSIS

#### School Screening

There have been several publications on screening for scoliosis since 2007, including a systematic review of



Figs 5A and B: (A) Girl with scoliosis; and (B) Adam's forward bend test



**Fig. 6:** Plumb line showing truncanal shift to right

the literature and large retrospective studies.<sup>39-41</sup> Labelle et al,<sup>39</sup> published a consensus statement developed by an international task force of the SRS regarding screening for AIS. The task force performed a systematic review of the literature through 2012 and used a modified Delphi process following the framework of the World Health Organization to reach consensus on the validity of a screening program. The panel reached consensus on the five domains studied, with four of the domains: Technical efficacy and clinical, program, and treatment effectiveness, supportive of screening, but there was insufficient evidence to make a statement with respect to cost-effectiveness.

The Adam's forward bend test with the use of a scoliometer (a specialized inclinometer) was agreed upon by the SRS task force as an effective quantitative measure, with 5 to 7° of deformity as a threshold for positive screening. The clinical effectiveness of screening for the detection of curves greater than 20° was supported in a large retrospective study by Luk et al.<sup>40</sup> Out of 115,190 adolescents followed until the age of 19 years. In their study, 2.8% of adolescents were referred for a radiograph. At final follow-up, the positive predictive value for spinal curvature greater than 20° was 43.8 and 9.8% for treatment. Sensitivity was near 90% for both diagnosis and treatment. Conversely, Yawn et al<sup>41</sup> reported on a population-based school screening program in Rochester, Minnesota. In this retrospective cohort study, 4.1% (92 out of 2,242) of the children screened positively and were referred for evaluation. The positive predictive value was low (0.05), and they concluded that roughly 450 children would need to be screened for every child who subsequently received treatment as a result of screening. The discrepancy in these studies points out the need

for effective screening systems, as inappropriate false-positive screening may lead to unnecessary referrals and radiographs with higher population cost. Although well-done population screening may be an effective means to capture all children at risk, many communities may not have sufficient resources to carry out these programs.

### Bracing

Treatment with rigid bracing (thoracolumbosacral and lumbosacral orthoses) is the most common nonoperative strategy to prevent curve progression.<sup>42-44</sup> Many different designs exist, but all attempt to restore the normal contours and alignment of the spine through the use of external forces and, with some designs, the stimulation of active correction as the patient moves the spine away from pressures within the brace. Permanent correction of the curve is typically not expected. Instead the brace functions as a holding device during the high-risk growth phase. Bracing is generally indicated for curves between 20 and 40° in adolescents who still have significant skeletal growth remaining (Risser status 0-2). The recommended wear time varies across clinicians, ranging from 12 to 23 hours per day until skeletal maturity is reached (2-4 years of treatment).<sup>45</sup>

In a hallmark study, Weinstein et al<sup>46</sup> reported on the efficacy of brace treatment bracing in adolescent idiopathic scoliosis trial (BrAIST). The study reported 75% success rate in patients randomly assigned to the bracing group as opposed to 42% in the group randomly assigned to observation alone. Success was defined as failure of progression of curve to 50° or more at skeletal maturity (Risser 4 for girls and Risser 5 for boys).

Katz et al,<sup>45</sup> demonstrated the efficacy of bracing in a non-controlled population, where 82% of patients who wore a brace for greater than 12 hours per day had less than 5° of curve progression compared to only 31% of those who wore the brace for less than 7 hours per day. An important feature of this study was that a temperature-sensitive data recorder imbedded in the spinal orthosis monitored brace wear compliance.

Bracing, however, has many disadvantages for patients including the need for radiographs to monitor brace fit and curve response, interference with sports and other activities, limited clothing choices, and self-consciousness about the brace. Brace wear for many patients is also a reminder of their medical condition.

### Surgical Treatment of Scoliosis

The accepted indications for surgery in adolescent scoliosis are curves greater than 50° with associated cosmetic implications. The goals of surgery include correction of deformity, prevent deterioration, and improve cosmesis.

Hibbs<sup>47</sup> reported early results from 59 patients, most with paralytic scoliosis, who had undergone posterior fusion surgery. His fusion technique involved elevating the bone flaps from the laminae and turning these flaps upward or downward to bring them into contact with the adjacent decorticated laminae.

Modern instrumentation surgery for scoliosis was pioneered in the early 1950s by Harrington,<sup>48</sup> who used a stainless steel rod-and-hook system to correct spinal deformities. Luque<sup>49</sup> reported a segmental spinal instrumentation method that used rods and sublaminar wiring. Unlike Harrington's system, which uses a distractive force applied to the spine through hooks, Luque's system used transverse forces applied segmentally through sublaminar wires.

Cotrel and Doubusset<sup>50</sup> described a multisegmental system, called CD instrumentation, in which multiple laminal and pedicle hooks and later, pedicle screws (PSs) were placed on the concave and convex sides of the curve. This allowed multiple fixation points using hooks, translation of the spine, and the creation of kyphosis by rod rotation. Cotrel and Doubusset believed that the CD system allowed for a shorter fusion area, derotation of the spine and the creation of kyphosis and lordosis in the thoracic spine and lumbar spine respectively.


Suk et al,<sup>51</sup> reported the first in a case series using PSs for fixation of the thoracic curve, and compared the results of three different surgical constructs: Hooks only, screws only, or a combination of screws and hooks. They found that PS constructs provided better correction of coronal, sagittal, and rotational deformity with less loss of correction, a shorter fusion area, and less

risk of neurological complications. In Japan in 1992, Abe et al also reported using PS constructs to correct thoracic curves in three patients, with a 78% correction rate. Several researchers have since compared surgical results between patients treated with PS-only constructs and those with hybrid constructs of hooks, sublaminar wires, and PSs. Kim et al,<sup>52</sup> compared the outcomes for patients treated with PS or hybrid constructs (29 each) and found that PS constructs offered better correction of the major curve and more improved pulmonary function than hybrid constructs, whereas the junctional change, lowest instrumented vertebra, time in surgery, and postoperative SRS-24 outcome scores were similar in both groups. Other authors echoed Kim's findings in retrospective comparative studies and systemic reviews, reporting that PS constructs provided better, or at least similar, correction and maintenance of the main curve, and required fewer revisions due to their biomechanical stability.<sup>53</sup>

The PS constructs have an important advantage in that vertebrae can be derotated directly with the PSs, thus reducing vertebral rotation by 42 to 60 % and reducing thoracic and lumbar hump.<sup>54</sup> The PS constructs also allow for osteotomies, including Ponte and pedicle-subtraction osteotomies and posterior vertebral column resection, so that even rigid and severe curves can be corrected efficiently without anterior procedures (Figs 7A to E).

### Anterior Approach Surgery

Dwyer developed an anterior instrumentation system using a titanium cable and screws to correct scoliosis.



**Figs 7A to E:** (A and B) Preoperative radiographs; (C) side bending radiograph; and (D and E) postoperative radiographs after Ponte osteotomies and pedicle screw instrumentation

After thorough discectomy and a morselized rib graft, the cable was threaded through the screw heads, and a tensioning device was applied to approximate the adjacent vertebral bodies. Fusion was achieved in 91% of 51 patients treated with this device; however, there was a loss of correction in 19 patients, and others experienced loss of lumbar lordosis and instrumentation failure.<sup>55</sup> Zielke et al,<sup>56</sup> developed ventral derotation spondylodesis (VDS), an anterior instrumentation system that was claimed to allow derotation and restoration of lordosis of the thoracolumbar spine, and to yield better correction than either the Harrington or Dwyer systems. After following 53 patients for at least 10 years after undergoing treatment with Dwyer or Zielke instrumentation, Otani<sup>57</sup> reported a 62% correction rate, a 6% rate of instrumentation failure, and patient satisfaction in most cases. However, other researchers have reported implant failure, loss of correction, progressive kyphosis, and pseudarthrosis in association with the VDS system. Kaneda et al<sup>58</sup> treated 25 patients with thoracolumbar or lumbar curves using an anterior dual-rod system, and obtained a correction rate of 83% for scoliosis and 86 % for rotation with restoration of lumbar lordosis. This 2-rod system is biomechanically robust enough to prevent loss of correction after surgery.

The main advantage of an anterior approach to the thoracic spine is preservation of mobile segments through a shorter fusion: In some scoliosis' curve patterns both proximal and distal levels can be saved, which would have been instrumented in a posterior spinal fusion. Other advantages of an anterior approach are resection of the growth plate and thereby prevention of the crankshaft phenomenon, and correction of the deformity by compression rather than distraction, which theoretically should have a lower rate of neurological complications.

## FUTURE ADVANCES

### Fusionless Surgery

Whereas conventional instrumented fusion for scoliosis has stood the test of time, it has the disadvantage of fusing mobile segments of the spine. Vertebral body stapling (VBS) utilizes C-shaped nitinol staples to compress across convex vertebral physes in growing children with IS with the goal of halting or reversing progressive curvature. The concept is to correct deformity without fusing the spine. The recommended indications are patients with Risser sign of 0 or 1 and coronal curve measuring between 20 and 45° deemed to be at risk of progression. Betz et al<sup>59</sup> using VBS in 41 patients followed for 2 years postoperatively reported success rate of 87% in all lumbar curves and in 79% of thoracic curves less than 35°.

## GENETIC TESTING

Genetic markers associated with AIS progression have the potential to be used as a prognostic test. Linkage analyses have identified candidate regions on chromosomes. A DNA-based prognostic test was developed by Ward et al,<sup>60</sup> to predict spinal curve progression in white patients with AIS. The test algorithm incorporates genotypes for 53 DNA markers and the patient's presenting Cobb's angle. The test reports a score between 1 and 200; this prognostic score correlates with the patient's risk of progression.

The results have not been replicated in other studies and there may be a variation based on ethnicity.<sup>61</sup>

## REFERENCES

1. Weinstein SL, Dolan LA, Cheng JC, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. *Lancet* 2008 May 3;371(9623):1527-1537.
2. Rogala EJ, Drummond DS, Gurr J. Scoliosis: incidence and natural history. A prospective epidemiological study. *J Bone Joint Surg Am* 1978 Mar;60(2):173-176.
3. Brooks HL, Azen SP, Gerberg E, Brooks R, Chan L. Scoliosis: A prospective epidemiological study. *J Bone Joint Surg Am* 1975 Oct;57(7): 968-972.
4. Bunnell WP. The natural history of idiopathic scoliosis. *Clin Orthop Relat Res* 1988 Apr;(229):20-25.
5. Kesling KL, Reinker KA. Scoliosis in twins. A meta-analysis of the literature and report of six cases. *Spine (Phila Pa 1976)* 1997 Sep 1;22(17):2009-2014; discussion 2015.
6. Garland HG. Hereditary scoliosis. *Br Med J* 1934 Feb 24; 1(3816):328.
7. Grauers A, Danielsson A, Karlsson M, Ohlin A, Gerdhem P. Family history and its association to curve size and treatment in 1,463 patients with idiopathic scoliosis. *Eur Spine J* 2013 Nov;22(11):2421-2426.
8. Thillard MJ. Vertebral column deformities following epiphysectomy in the chick. *C R Hebd Seances Acad Sci* 1959 Feb;248(8): 1238-1240.
9. Machida M, Dubousset J, Imamura Y, Iwaya T, Yamada T, Kimura J. Role of melatonin deficiency in the development of scoliosis in pinealectomized chickens. *J Bone Joint Surg Br* 1995 Jan;77(1):134-138.
10. Machida M, Saito M, Dubousset J, Yamada T, Kimura J, Shibasaki K. Pathological mechanism of idiopathic scoliosis: experimental scoliosis in pinealectomized rats. *Eur Spine J* 2005 Nov;14(9):843-848.
11. Cheung KM, Wang T, Poon AM, Carl A, Tranmer B, Hu Y, Luk KD, Leong JC . The effect of pinealectomy on scoliosis development in young nonhuman primates. *Spine (Phila Pa 1976)* 2005 Sep 15;30(18):2009-2013.
12. Bagnall KM, Raso VJ, Hill DL, Moreau M, Mahood JK, Jiang H, Russell G, Bering M, Buzzell GR. Melatonin levels in idiopathic scoliosis. Diurnal and nocturnal serum melatonin levels in girls with adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 1996 Sep 1;21(17):1974-1978.
13. Lowe T, Lawellin D, Smith D, Price C, Haheer T, Merola A, O'Brien M. Platelet calmodulin levels in adolescent idiopathic scoliosis: do the levels correlate with curve progression and severity? *Spine (Phila Pa 1976)* 2002 Apr 1;27(7):768-775.

14. Dickson RA, Lawton JO, Archer IA, Butt WP. The pathogenesis of idiopathic scoliosis. Biplanar spinal asymmetry. *J Bone Joint Surg Br* 1984 Jan;66(1):8-15.
15. Herman R, MacEwen D. Idiopathic scoliosis: a visio-vestibular disorder of the central nervous system. In: Zorab PA, editor. *Scoliosis. Proceedings of the sixth symposium*. Academic Press; 1979. p. 61-69.
16. Chu WC, Lam WW, Chan YL, Ng BK, Lam TP, Lee KM, Guo X, Cheng JC. Relative shortening and functional tethering of spinal cord in adolescent idiopathic scoliosis?: study with multiplanar reformat magnetic resonance imaging and somatosensory evoked potential. *Spine (Phila Pa 1976)* 2006 Jan 1;31(1):E19-25.
17. Abul-Kasim K, Overgaard A, Karlsson MK, Ohlin A. Tonsillar ectopia in idiopathic scoliosis: does it play a role in the pathogenesis and prognosis or is it only an incidental finding? *Scoliosis* 2009 Nov;4:25.
18. Goldberg CJ, Dowling FE, Fogarty EE Adolescent idiopathic scoliosis—early menarche, normal growth. *Spine (Phila Pa 1976)* 1993 Apr; 18(5):529-535.
19. Lonstein JE, Carlson JM. The prediction of curve progression in untreated idiopathic scoliosis during growth. *J Bone Joint Surg Am* 1984 Sep;66(7):1061-1071.
20. Risser JC. The classic: The iliac apophysis; an invaluable sign in the management of scoliosis. 1958. *Clin Orthop Relat Res* 2010 Mar; 468(3):643-653.
21. Bitan FD, Veliskakis KP, Campbell BC. Differences in the Risser grading systems in the United States and France. *Clin Orthop Relat Res* 2005 Jul;(436):190-195.
22. Ascani E, Bartolozzi P, Logroscino CA, Marchetti PG, Ponte A, Savini R, Travaglini F, Binazzi R, Di Silvestre M. Natural history of untreated idiopathic scoliosis after skeletal maturity. *Spine (Phila Pa 1976)*;1986 Oct;11(8):784-789.
23. Weinstein SL. Idiopathic scoliosis. Natural history. *Spine (Phila Pa 1976)* 1986 Oct;11(8):780-783.
24. Sanders JO, Khoury J, Kishan S Browne RH, Mooney JF 3rd, Arnold KD, McConnell SJ, Bauman JA, Finegold DN. Predicting scoliosis progression from skeletal maturity: a simplified classification during adolescence. *J Bone Joint Surg Am* 2008 Mar; 90(3):540-553
25. Frymoyer JW, Pope MH, Clements JH, Wilder DG, McPherson B, Ashikaga T. Risk factors in low-back pain. An epidemiological survey. *J Bone Joint Surg Am* 1983 Feb; 65(2): 213-218
26. Balague F, Skovron ML, Nordin M, Dutoti G, Pol LR, Waldburger M. Low back pain in schoolchildren. A study of familial and psychological factors. *Spine (Phila Pa 1976)* 1995 Jun 1; 20(11):1265-1270
27. Burton AK, Clarke RD, McClune TD, Tillotson KM The natural history of low back pain in adolescents. *Spine (Phila Pa 1976)* 1996 Oct 15; 21(20):2323-2328.
28. Ramirez N, Johnston CE, Browne RH. The prevalence of back pain in children who have idiopathic scoliosis *J Bone Joint Surg Am* 1997 Mar;79(3):364-368.
29. Sato T; Hirano T; Ito T; Morita O; Kikuchi R, Endo N, Tanabe N. Back pain in adolescents with idiopathic scoliosis: epidemiological study for 43,630 pupils in Niigata City, Japan. *Eur Spine J* 2011 Feb;20 (2): 274-279
30. Gazioglu K, Goldstein LA, Femi-Pearse D, Yu PN. Pulmonary function in idiopathic scoliosis. Comparative evaluation before and after orthopaedic correction. *J Bone Joint Surg Am* 1968 Oct;50(7):1391-1399.
31. Makley JT, Herndon CH, Inkley S, Doershuk C, Matthews LW, Post RH, Littell AS. Pulmonary function in paralytic and non-paralytic scoliosis before and after treatment. A study of sixty-three cases. *J Bone Joint Surg Am* 1968 Oct;50(7):1379-1390.
32. Upadhyay SS, Mullaji AB, Luk KD, Leong JC. Evaluation of deformities and pulmonary function in adolescent idiopathic right thoracic scoliosis. *Eur Spine J* 1995;4(5):274-279.
33. Kumano K, Tsuyama N. Pulmonary function before and after surgical correction of scoliosis. *J Bone Joint Surg Am* 1982 Feb;64(2):242-248.
34. Weinstein SL, Zavala DC, Ponseti IV. Idiopathic scoliosis: long-term follow-up and prognosis in untreated patients. *J Bone Joint Surg Am* 1981 Jun;63(5):702-712.
35. Kim YJ, Lenke LG, Bridwell KH, Kim KL, Steger-May K. Pulmonary function in adolescent idiopathic scoliosis relative to the surgical procedure. *J Bone Joint Surg Am* 2005 Jul;87(7):1534-1541.
36. King, HA, Moe JH, Bradford DS, Winter RB. The selection of fusion levels in thoracic idiopathic scoliosis. *J Bone Joint Surg Am* 1983 Dec; 65(9): 1302-1313.
37. Lenke LG, Betz R, Harms J, Bridwell KH, Clements DH, Lowe TG, Blanke K. Adolescent idiopathic scoliosis: a new classification to determine extent of spinal arthrodesis. *J Bone Joint Surg Am* 2001 Aug; 83(8):1169-1181.
38. Cummings RJ, Loveless EA, Campbell J, Samelson S, Mazur JM. Interobserver reliability and intraobserver reproducibility of the system of King et al. for the classification of adolescent idiopathic scoliosis. *J Bone Joint Surg Am* 1998 Aug;80(8):1107-1111.
39. Labelle H, Richards SB, De Kleuver M, Grivas TB, Luk KD, Wong HK, Thometz J, Beauséjour M, Turgeon I, Fong DY: Screening for adolescent idiopathic scoliosis: an information statement by the scoliosis research society international task force. *Scoliosis* 2013 Oct 31;8:17.
40. Luk KD, Lee CF, Cheung KM, Cheng JC, Ng BK, Lam TP, Mak KH, Yip PS, Fong DY. Clinical effectiveness of school screening for adolescent idiopathic scoliosis: a large population-based retrospective cohort study. *Spine (Phila Pa 1976 )* 2010 Aug 1;35(17):1607-1614.
41. Yawn BP, Yawn RA. The estimated cost of school scoliosis screening. *Spine (Phila Pa 1976 )* 2000 Sep 15;25(18):2387-2391.
42. Emans JB, Kaelin A, Bancel P, Hall JE, Miller ME. The Boston bracing system for idiopathic scoliosis. Follow-up results in 295 patients. *Spine. (Phila Pa 1976)* 1986 Oct;11(8):792-801.
43. Price CT, Scott DS, Reed FE Jr, Riddick MF. Nighttime bracing for adolescent idiopathic scoliosis with the Charleston bending brace. Preliminary report. *Spine (Phila Pa 1976)* 1990 Dec;15(12):1294-1299.
44. Trivedi JM, Thomson JD. Results of Charleston bracing in skeletally immature patients with idiopathic scoliosis. *J Pediatr Orthop* 2001 May-Jun;21(3):277-280.
45. Katz DE, Herring JA, Browne RH, Kelly DM, Birch JG. Brace wear control of curve progression in adolescent idiopathic scoliosis. *J Bone Joint Surg Am* 2010 Jun;92(6):1343-1352.
46. Weinstein SL, Dolan LA, Wright JG, Dobbs MB. Effects of bracing in adolescents with scoliosis. *N Eng J Med* 2013 Oct 17;369(16):1512-1521.
47. Hibbs RA. A report of fifty-nine cases of scoliosis treated by the fusion operation. *J Bone Joint Surg Am* 1924 Jan;6(1):3-34.
48. Harrington PR. Treatment of scoliosis: correction and internal fixation by spine instrumentation. *J Bone Joint Surg Am* 1962 Jun;44-A:591-610.

49. Luque ER. Segmental spinal instrumentation for correction of scoliosis. *Clin Orthop Relat Res* 1982 Mar;(163):192-8.
50. Cotrel Y, Dubousset J, Guillaumat M. New universal instrumentation in spinal surgery. *Clin Orthop Relat Res* 1988 Feb;227: 10-23.
51. Suk SI, Lee CK, Min HJ, Cho KH, Oh JH. Comparison of Cotrel-Dubousset pedicle screws and hooks in the treatment of idiopathic scoliosis. *Int Orthop* 1994;18(6):341-346.
52. Kim YJ, Lenke LG, Kim J, Bridwell KH, Cho SK, Cheh G, Sides B. Comparative analysis of pedicle screw versus hybrid instrumentation in posterior spinal fusion of adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2006 Feb 1;31(3):291-298.
53. Ledonio CG, Polly DW Jr, Vitale MG, Wang Q, Richards BS. Pediatric pedicle screws: comparative effectiveness and safety: a systematic literature review from the Scoliosis Research Society and the Pediatric Orthopaedic Society of North America task force. *J Bone Joint Surg Am* 2011 Jun 6;93(13): 1227-1234.
54. Lee SM, Suk SI, Chung ER. Direct vertebral rotation: a new technique of three-dimensional deformity correction with segmental pedicle screw fixation in adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2004 Feb 1;29(3):343-349.
55. Dwyer AF, Schafer MF. Anterior approach to scoliosis. Results of treatment in fifty-one cases. *J Bone Joint Surg Br* 1974 May;56(2):218-224.
56. Zielke K, Stunkat R, Beaujean F. [Ventralspondylodesis (author's transl)]. *Arch Orthop Unfallchir* 1976 Aug 19;85(3):257-77.
57. Otani K, Saito M, Sibasaki K. Anterior instrumentation in idiopathic scoliosis: a minimum follow-up of 10 years. *Int Orthop* 1997;21(1):4-8.
58. Kaneda K, Shono Y, Satoh S, Abumi K. New anterior instrumentation for the management of thoracolumbar and lumbar scoliosis. Application of the Kanada two-rod system. *Spine (Phila Pa 1976)* 1996 May 15;21(10):1250-1261.
59. Betz RR, Ranade A, Samdani AF, Chafetz R, D'Andrea LP, Gaughan JP, Asghar J, Grewal H, Mulcahey MJ. Vertebral body stapling: a fusionless treatment option for a growing child with moderate idiopathic scoliosis. *Spine (Phila Pa 1976)* 2010 Jan 15;35(2):169-1676.
60. Ward K, Ogilvie JW, Singleton MV, Chettier R, Engler G, Nelson LM. Validation of DNA-based prognostic testing to predict spinal curve progression in adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2010 Dec 1;35(25):E1455-E1464.
61. Ogura Y, Takahashi Y, Kou I, Nakajima M, Kono K, Kawakami N, Uno K, Ito M, Minami S, Yanagida H. A replication study for association of 53 single nucleotide polymorphisms in a scoliosis prognostic test with progression of adolescent idiopathic scoliosis in Japanese. *Spine (Phila Pa 1976)* 2013 Jul 15;38(16):1375-1379.