

Clinical Profile of Refractive Errors in School-going Children of Bareilly, Uttar Pradesh, India

¹Juhi Mittal, ²Yusuf Rizvi, ³Ritu Jain

ABSTRACT

Introduction: Refractive error is one of the important causes of impaired visual acuity in children.

Aim: To study the pattern of refractive errors in children of Bareilly city and determine its association with selected variables of age, sex, educational status, socioeconomic status, residing locality, family history, previous use of spectacles, amblyopia, and strabismus.

Materials and methods: A cross-sectional study was conducted on 521 children aged 5 to 15 years attending the ophthalmology outpatient department (OPD) of a teaching hospital of Bareilly. Visual status along with dry and cycloplegic refraction was carried out. Relevant data in relation to the selected variables were compiled. Statistical analysis of obtained results was carried out using the software Statistical Package for the Social Sciences (SPSS) version 22.

Results: The mean age of presentation of refractive error was 10.75 ± 2.96 years with a distinct male preponderance and with a male:female ratio of 1.33:1. Parental education background revealed fair literacy rate with 56.23% mothers and 69.28% fathers having intermediate qualification and above. Higher incidence of refractive error was noted in children with upper lower socioeconomic status (34.54%) and with those having positive family history (61.23%); 7.67% children were amblyopic and 7.10% children had coexisting strabismus. Most of the children (80.23%) gave no history of previous use of spectacles. Astigmatism was noted as the commonest refractive error in 45.09% followed by myopia (42.86%) and hypermetropia (12.05%).

Conclusion: The study provides an insight into the quantum, extent, and form of visual impairment prevalent in Bareilly district. It furnishes essential data for planning and evaluating preventive and curative services for visual impairment of children in this region.

Clinical significance: These data support the assumption that vision screening of school children in developing countries would be very useful in early detection of correctable causes of poor vision, especially refractive errors and in preventing visual complications.

Keywords: Refractive errors, School-going children.

How to cite this article: Mittal J, Rizvi Y, Jain R. Clinical Profile of Refractive Errors in School-going Children of Bareilly, Uttar Pradesh, India. *Int J Adv Integ Med Sci* 2017;2(4):169-172.

¹Junior Resident (Third Year), ^{2,3}Associate Professor

¹⁻³Department of Ophthalmology, Rohilkhand Medical College & Hospital, Bareilly, Uttar Pradesh, India

Corresponding Author: Juhi Mittal, Junior Resident (Third Year), Department of Ophthalmology, Rohilkhand Medical College & Hospital, Bareilly, Uttar Pradesh, India, Phone: +919899696851, e-mail: dr.juhimittal@gmail.com

Source of support: Nil

Conflict of interest: None

INTRODUCTION

Refractive error is one of the most common causes of visual impairment around the world and the second leading cause of treatable blindness.¹

The significance of early detection of refractive errors in childhood springs from the fact that this condition is largely avoidable with a cost-effective treatment.² Screening activity is essential since children do not complain of defective vision, and at times may be unaware of their problem. It is noteworthy that poor vision during childhood and adolescence affects academic and overall school performance and may have a negative influence on the future life. Available data on the type of refractive errors in developing countries are scarce. It shows interregional disparities based on geography, urban rural background, and ethnicity.

Keeping these factors in view, the present study was undertaken to determine the prevalence of refractive errors in children of Bareilly district.

MATERIALS AND METHODS

This study was done on school-going children aged 5 to 15 years, attending the ophthalmology OPD of a tertiary care center in Bareilly city. A prior informed consent was taken from the parents of participating children. A detailed history was taken about the present and past ocular problems along with the history of use of spectacles. Both unaided and aided visual acuity were recorded using Snellen's chart. Postmydriatic refraction was done after 1 week of cycloplegic retinoscopy and appropriate spectacles were prescribed to the children as per the protocol. Spherical equivalent (SE) was used for calculations of refractive error. The SE was derived by adding the spherical component of refraction to half of the cylindrical component. Myopia was defined as an SE of at least -0.5 D and hyperopia as that of $+2.00$ D or more. Astigmatic students were labelled with a cylindrical refractive power of 0.75 D or more in at least one eye. Anisometropia was defined as a difference in SE of at least 1.0 D between the two eyes. Data were analyzed using SPSS, version 22 software program. Percentage and 95% CI were used to describe the prevalence of

refractive errors. Spearman chi-squared test was applied for qualitative data. A p-value less than 0.05 was considered statistically significant.

RESULTS

A total of 521 children were studied over a period of 1 year. Out of these, 13 children had unilateral refractive error, hence 1,029 eyes with refractive errors of the 521 children were taken into account. Out of these, 57.2% were boys. A total of 235 children (45.11%) were in the age range of 5 to 10 years and the remaining 286 (54.89%) were 11 to 15 years. In the present study, the average age of presentation of children with refractive error was 10.75 ± 2.96 years.

It was observed that 298 (57.20%) were from urban areas, while 223 (42.80%) children were from rural areas (Table 1).

This study demonstrated that 293 (56.23%) mothers and 361 (69.28%) fathers were having educational qualification of intermediate and above (Table 2).

It was observed that visual acuity at the time of presentation was better than 6/12 in 46.55% of eyes, 6/18 to 6/36 in 34.49% of eyes, and ≤6/60 in 18.95% of the eyes. Hence, most of the children had mild-to-moderate visual impairment.

In terms of socioeconomic status, 68 (13%) of the children belonged to families of upper class, 71 (13.63%) belonged to upper middle class, 159 (30.51%) to lower middle class, 180 (34.54%) to upper lower class, and only 43 (8.25%) children belonged to lower class families. Hence, majority of children, both male and female, were noted to come from upper lower-class families.

In the present study, 319 (61.23%) children were found to have a positive family history of refractive errors

Table 1: Distribution of children with refractive errors based on rural–urban background (n = 521)

Area	Male	Female	Total
Urban	167 (56.05%)	131 (43.95%)	298 (57.20%)
Rural	131 (58.75%)	92 (41.25%)	223 (42.80%)
Total	298 (57.20%)	223 (42.80%)	521 (100%)

$\chi_{(2)}^2 = 2.769; p = 0.5914$

Table 2: Distribution of children with refractive errors based on educational status of parents (n = 521)

Education	Mother	Father
Illiterate	49 (9.40%)	22 (4.22%)
Primary	95 (18.23%)	35 (6.74%)
High school	84 (16.13%)	103 (19.76%)
Intermediate	128 (24.57%)	131 (25.14%)
Graduate and above	165 (31.67%)	230 (44.14%)
Total	521 (100%)	521 (100%)

$\chi_{(10)}^2 = 21.82; p = 1.00$

Table 3: Distribution of children with refractive errors based on family history of refractive errors (n = 521)

Family history of refractive error	Male	Female	Total
Present	179 (60.06%)	140 (62.78%)	319 (61.23%)
Absent	119 (39.93%)	83 (37.22%)	202 (38.77%)
Total	298 (57.2%)	223 (42.80%)	521 (100%)

$\chi_{(1)}^2 = 14.586; p = 0.5857$

Table 4: Distribution of children with refractive errors based on history of previous use of spectacles (n = 521)

H/O previous use of spectacles	Male	Female	Total
Present	58 (56.31%)	45 (43.69%)	103 (19.77%)
Absent	240 (57.42%)	178 (42.58%)	418 (80.23%)
Total	298 (57.30%)	223 (42.70%)	521 (100%)

$\chi_{(2)}^2 = 3.876; p = 0.4458$

with 202 (38.77%) children reported no such history. Association of refractive errors was hence not seen to be significantly associated with a positive family history ($p > 0.05$) (Table 3).

In this study, 40 (7.67%) children were detected to have amblyopia, out of which anisometropic amblyopia was present in 24 (60%) and strabismic amblyopia in 16 (40%) children.

Out of these 521 children, 37 (7.10%) children had strabismus. Esotropia was the most common type of strabismus in children with refractive errors, accounting for 64.70%, while exotropia was present in 35.30% children.

In our study, only 103 (19.77%) children were already using spectacles, of which 58 (56.31%) were males and 45 (43.69%) were females. However, majority of the children (80.23%) with refractive errors were not using spectacles previously. There was no significant difference between males and females using spectacles (Table 4).

In the current study, the most common type of refractive error was astigmatism (45.09%), followed by myopia (42.86%) and hypermetropia (12.05%). No association between the type of refractive error and laterality of eye was noted. An age-related shift was observed from hypermetropia with a higher incidence in younger age group to myopia and astigmatism with a higher incidence in older age group (Table 5).

Table 5: Distribution of types of refractive error by age (n = 1029)

Type of refractive error	5–10 years	11–15 years	Total
Myopia	203 (46.03%)	238 (53.96%)	441 (42.86%)
Hypermetropia	63 (50.81%)	61 (49.19%)	124 (12.05%)
Astigmatism	203 (43.75%)	261 (56.25%)	464 (45.09%)
Total	469 (45.57%)	560 (54.43%)	1029 (100%)

$\chi_{(2)}^2 = 15.353; p = 0.7190$



DISCUSSION

Childhood blindness due to uncorrected refractive error has emerged as a major public health problem, the cognizance of which has been taken by the World Health Organization in its Vision 2020 program. Worldwide, uncorrected refractive error accounts for up to 42% of visual impairment and equally affects both developing and developed nations.³ Interregional disparities in the nature, demography, and type of refractive error have been reported by various studies. A close analysis of these is particularly useful in educating scientific world and public for effective management of the condition.

The present study has been designed as a cross-sectional hospital-based descriptive study on school-aged children falling in the age group of 5 to 15 years hailing largely from the Bareilly district of northern India.

Of the 521 assessed children during the study period of 1 year, the mean age of presentation was noted to be 10.75 ± 2.96 years (5–15 years). This was comparable with reports of 9.3 ± 3.4 , 9.7 ± 3.3 , and 10.7 ± 3.1 years by Kalikivayi et al⁴ in Hyderabad, Wu et al⁵ in China, and Yamamah et al⁶ in Egypt respectively. The age of presentation of refractive error assumes large significance, since it determines the prognosis of permanent visual disabilities like amblyopia, strabismus and has a bearing on the intellect and overall psychological development of the child.

The gender distribution of refractive errors in our study revealed a higher preponderance of males, with 298 (57.20%) male and 223 (42.80%) female participants. This difference was, however, not statistically significant ($p > 0.05$). A hospital-based study by Rai et al⁷ in Nepal reported similar results with 58% male affliction. Comparable results were reported for other hospital-based studies done by Alam and Fareed⁸ and Sethi et al⁹ in Pakistan, Matta et al¹⁰ in New Delhi, and Rohul et al¹¹ in Kashmir.

In our study, myopia was noted in 42.86% of children while astigmatism and hypermetropia in were noted in 45.09 and 12.05% of patients respectively. This is in agreement with the study done by Sethi et al¹² in Pakistan and similar studies in Nepal,⁷ Ethiopia,¹³ and Egypt,¹⁴ all of which designated astigmatism as the most common type of refractive error followed by myopia and hypermetropia.

The association of refractive error with hereditary factors has been acknowledged by most studies. Our study noted a parental or sibling background of refractive error in 61.23% of patients. It was not significant in our study, but Pavithra et al¹⁵ found a very strong relationship between refractive errors and hereditary or familial factors. Ali et al¹⁶ also reported that a positive family history of myopia is related to progression of myopia and refractive error.

The prevalence of refractive error was found to be high in the urban area (57.20%) as compared with rural areas (42.80%), similar to the findings of Pavithra et al¹⁵ in Bengaluru, Batra et al¹⁷ in Punjab, and Padhye et al² in Maharashtra. Comparatively lower rates of refractive error are seen in rural areas where children do not face the same emphasis on schooling and are frequently withdrawn from school at an early age, whereas the children of urban areas have better access to educational services which make the children more prone for near work compared with rural children.

It was seen that parents (both mother and father) with higher levels of education were more likely to have children with refractive errors. Similar results of association between educational background and refractive errors in children were observed in the New Delhi survey.¹⁸ Although these studies did not consider the effect of mother's educational status on the refractive state of child, it was found significant in our study.

Refractive error is one of the few ailments that equally affect the affluent and the poor, the developing and the developed world. Socioeconomic status seems to be an indicator determining the type and degree of refractive error. Contemporary studies have documented a high association of myopia and other refractive error with high socioeconomic background and better schooling access. Our study noted a high association of refractive error with middle-income background with more than 60% affected children coming from either lower middle or upper lower class (refer Kuppaswamy classification for urban and BJ Prasad's classification for rural economic stratification).

In the present study, only 103 out of 521 children with refractive errors were using spectacles (19.77%). The percentage of such children was notably much higher (57%) in a similar study done in Nepal.⁷ The reported differences highlight the indifference of parents to the need of wearing spectacles by needy children of the area. A possible reason could be a social stigma with glasses especially for girls, lack of facilities of spectacle provisioning, or simply the irresponsible attitude toward children at large. Counseling of parents and societal awareness are therefore of equal importance as that of diagnosing refractive errors and prescribing glasses in this scenario.

A total of 40 (7.67%) children were found to have amblyopia. Similar results were seen in the studies done by Pant et al¹⁹ in Nepal, and Sethi et al¹² in which 7.62 and 6.95% children were amblyopic respectively. Amblyopia treatment is most effective when done early in a child's life, usually before the age of 7 years.¹⁵ This reinforces the need to screen for amblyopia in all children presenting with refractive error.

In this study, out of 521 children who were studied, 37 (7.10%) were found strabismic. A study done by

Kalikivayi et al⁴ in Southern India demonstrated that 13.3% children with refractive errors had strabismus, but the association of strabismus with refractive errors was not found to be significant. A higher prevalence was noted in Pakistan by Sethi et al,¹² where 20% of children with refractive errors had strabismus. This difference may partly be due to the inclusion of much younger age group in the study.

Present-day studies investigating refractive errors in children are essentially population-based surveys. They employ identical protocols as outlined by the refractive error survey in children employing cluster sampling. To that extent, our study carries the handicap of a hospital-based study confined to eye OPD of a tertiary hospital. This provided the data with a distinct urban bias, since majority of rural children failed to report particularly with milder degrees of visual impairment. Despite these limitations, the study provided an insight into the quantum, extent, and form of visual impairment prevalent in Bareilly district. It furnishes essential data for planning and evaluating preventive and curative services for visual impairment of children in this region.

CONCLUSION

In summary, the average age of presentation of refractive errors in children was 10.90 ± 3.16 years. A positive family history and higher education of parents was significantly associated with the presence of refractive error in child. Majority of children with refractive errors belonged to lower middle-class families. Amblyopia and strabismus were also noted in a few children with refractive errors. Most of the children with refractive error present with mild-to-moderate decrease in visual acuity ($\leq 6/36$). Astigmatism was the most common type of refractive error followed by myopia and hypermetropia. Most of the children suffered from mild-to-moderate degree of refractive error in all categories. An age-related shift from hypermetropia in younger age group to myopia in older age group was found. No significant association was found between age of children and the prevalence of astigmatism.

ACKNOWLEDGMENT

Authors would extremely like to thank the optometry staff for their help and cooperation.

REFERENCES

1. Resnikoff S, Pascolini D, Mariotti SP, Pokharel GP. Global magnitude of visual impairment caused by uncorrected refractive errors in 2004. *Bull World Health Organ* 2008 Jan;86(1):63-70.
2. Padhye AS, Khandekar R, Dharmadhikari S, Dole K, Gogate P, Deshpande M. Prevalence of uncorrected refractive error and other eye problems among urban and rural school children. *Middle East Afr J Ophthalmol* 2009 Apr-Jun;16(2):69-74.
3. Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol* 2012 May;96(5):614-618.
4. Kalikivayi V, Naduvilath TJ, Bansal AK, Dandona L. Visual impairment in school children in southern India. *Indian J Ophthalmol* 1997 Jun;45(2):129-134.
5. Wu JF, Bi HS, Wang SM, Hu YY, Wu H, Sun W, Lu TL, Wang XR, Jonas JB. Refractive error, visual acuity and causes of vision loss in children in Shangdon, China. The Shangdon Children Eye Study. *PLoS One* 2013 Dec;8(12):e82763.
6. Yamamah GA, Talaat Abdel Alim AA, Mostafa YS, Ahmed RA, Mahmoud AM. Prevalence of visual impairment and refractive errors in children of South Sinai, Egypt. *Ophthalmic Epidemiol* 2015 Aug;22(4):246-252.
7. Rai S, Thapa HB, Sharma MK, Dhakhwa K, Karki R. The distribution of refractive errors among children attending Lumbini Eye Institute, Nepal. *Nepal J Ophthalmol* 2012 Jan-Jun;4(7):90-95.
8. Alam M, Fareed M. Refractive errors; profile in school age children. *Professional Med J* 2011 Oct-Dec;18(4):649-653.
9. Sethi MJ, Sethi S, Iqbal R. Frequency of refractive error in children visiting outpatient department agency Headquarter Hospital Landi Kotil. *Gomal J Med Sci* 2009 Jul-Dec;7(2): 114-117.
10. Matta S, Matta P, Gupta V, Dev V. Refractive errors among adolescents attending ophthalmology OPD. *Indian J Community Med* 2006 Apr-Jun;31(2):114-115.
11. Rohul J, Maqbool A, Hussain SA, Shamila H, Anjum F, Hamdani ZA. Prevalence of refractive errors in adolescents in out-patient attendees of the preventive ophthalmology clinic of community medicine, S K I M S, Kashmir, India. *NUJHS* 2013 Mar;3(1):17-20.
12. Sethi S, Sethi MJ, Hussain I, Khan T. Refractive errors in children attending out-patient department of ophthalmology, Khyber Teaching Hospital, Peshawar. *Pak J Ophthalmol* 2008 Dec;4:1-5.
13. Mehari ZA. Pattern of childhood ocular morbidity in rural eye hospital, Central Ethiopia. *BMC Ophthalmol* 2014 Apr;14(1):50.
14. Shrestha GS, Sujakhu D, Joshi P. Refractive error among school children in Jhapa, Nepal. *J Optom* 2011 Apr;4(2):49-55.
15. Pavithra MB, Maheshwaran R, Rani Sujatha MA. A study on the prevalence of refractive errors among school children of 7-15 years age group in the field practice areas of a medical college in Bangalore. *Int J Med Sci Public Health* 2013 Jan;2(3):641-645.
16. Ali A, Ahmad I, Ayub S. Prevalence of undetected refractive errors among school children. *E:/Biomedica* 2007 Jul-Dec;23(Bio 21):96-101.
17. Batra N, Kaushal D, Gill AS. Refractive errors in school children: a review from Punjab. *NPCB India* 2007 Oct-Dec;1(4):2-4.
18. Murthy GV, Gupta SK, Ellwein LB, Munoz SR, Pokharel GP, Sanga L, Bachani D. Refractive error in children in an urban population in New Delhi. *Invest Ophthalmol Vis Sci* 2002 Mar;43(3):623-631.
19. Pant BP, Ghising R, Awasthi S, Pant SR, Bhatta RC. Refractive status among the students presenting to Geta Eye Hospital, Kailali, Nepal. *Nepal Med Coll J* 2010 Jun;12(2):95-99.