

# Prevalence and Factors Associated with Modic Changes of Lumbosacral Spine in Nepalese Patients with Chronic Low Back Pain

<sup>1</sup>Pankaj R Nepal, <sup>2</sup>Suman Rijal

## ABSTRACT

**Aim:** To determine prevalence and factors affecting MC of vertebral endplate in patients with chronic low back pain Nepalese patients coming to a neurosurgical–tertiary care center in Nepal.

**Materials and methods:**

**Study design:** Prospective analytical study

**Sample size:** 194 cases

**Sampling technique:** Non- probability consecutive sampling

**Data collection and analysis:** Patients clinical data like age, gender, occupation, body mass index, smoking habit, and history of diabetes mellitus (DM) were noted and status of Modic changes were noted from the MRI.

The analysis was done using SPSS-20. Mean, and standard deviation (SD) were calculated for quantitative variables. Frequency and percentage were calculated for qualitative variables. Effect modifier has been controlled through stratification and post-stratification Chi-square test was applied with  $P \leq 0.05$  was taken significantly.

**Results:** Overall prevalence of the Modic changes was 50.5% where MC II was the commonest of all the types. Prevalence of MC was significantly higher in patients who are elderly, obese, had severe pain on presentation, and had a habit of smoking.

**Conclusion:** Modic change occurring at the vertebral endplate is a common phenomenon with significant association with the age, severity of pain, smoking, and Body Mass Index (BMI).

**Clinical Significance:** Modic changes (MC) are the pathological changes of the vertebral body and endplate of the vertebra. Its prevalence in the general population is about 6% and presented in about 35% of patients with low back pain. Various researches have been conducted to see its association with back pain; however, controversies still persist. So this research further highlights the factors that could be associated with its prevalence.

**Keywords:** Low Back Ache, Lower Back Pain, Modic Changes, Herniated Lumbar Disc

**How to cite this article:** Nepal PR, Rijal S. Prevalence and Factors Associated with Modic Changes of Lumbosacral Spine in Nepalese Patients with Chronic Low Back Pain. *J Spinal Surg* 2018;5(3):115-119.

**Source of support:** Nil

**Conflict of interest:** None

## INTRODUCTION

Modic change (MC) was first described in MRI by de Roos et al. in 1987 as pathological changes of the vertebral body and endplate of the vertebra.<sup>1</sup> In 1986 its classification was formulated by Modic et al., into three types secondary to marrow edema, fatty degeneration and bony sclerosis (Table 1).<sup>2,3</sup>

As MC is primarily diagnosed by MRI,<sup>4</sup> its actual prevalence in general population is difficult to calculate; however, is quoted around 6% and is present in about 35% of patients with low back pain (LBP).<sup>5</sup> Since Modic et al. first described these changes, there have been various researches conducted to see its association with back pain; however, controversies still persist. MC Type I (MC-I) has a poor outcome in the patient with persistent LBP<sup>6</sup> and has less improvement of back pain following microdiscectomy if they have type I changes preoperatively.<sup>7</sup> These vertebral endplate changes are also the essential features of associated disease progression.<sup>8</sup>

Despite all these facts, controversies still persist. Hutton MJ et al. claimed that these vertebral endplate changes are reversible.<sup>4</sup> Similarly, Keller A et al. stated that they do not influence the clinical course of back pain,<sup>9</sup> and Jensen RK et al. showed their association with LBP are not constant.<sup>10</sup>

**Table 1:** Classification of Modic Changes.

Types	Underlying Pathology	T1	T2
MC- I	Marrow edema	Hypointense	Hyperintense
MC- II	Fatty degeneration	Hyperintense	iso-hyperintense
MC- III	Bony sclerosis	Hypointense	Hypointense

<sup>1</sup>Head of Department, <sup>2</sup>Consultant Neurosurgeon

<sup>1</sup>Department of Neurosurgery, BNC Medical College Teaching Hospital, Birtamode, Jhapa, State 1, Nepal

<sup>2</sup>Department of Neurosurgery, National Institute of Neurological and Allied Sciences, Kathmandu, State 3, Nepal

**Corresponding Author:** Pankaj R Nepal, Head of Department, Department of Neurosurgery, BNC Medical College Teaching Hospital, Birtamode, Jhapa, State 1, Nepal, Email: drprnepal@yahoo.com

With the objective to determine prevalence and factors affecting Modic changes of Lumbosacral vertebral endplate in patients with chronic low back pain, this study was conducted in a neurosurgical –tertiary care center in Nepal.

## MATERIALS AND METHODS

*Study design:* Prospective analytical study

*Sample size:* 194 cases

*Sampling technique:* Non-probability consecutive sampling

*Duration:* 3 months

*Inclusion criteria:* All male and female patients of 30 – 70 years of age, presented with moderate to severe pain, visual analogue scale (VAS) >3, for more than 3 months duration in this institute.

### Exclusion criteria

Patients presented with recent trauma

Obvious vertebral body injury or spinal injury

Obvious vertebral body or disc lesions like Metastasis or Tuberculosis

Obvious spinal deformity like scoliosis.

*Data collection:* All patients meeting the inclusion criteria were enrolled in the study. Patients clinical data like age, gender, occupation, body mass index, smoking habit, and history of diabetes mellitus (DM) were noted, and status of Modic changes was noted from the MRI of the patient (Fig. 1-3) based on its classification (Table 1).

### Data analysis

Analysis has been done through SPSS software; version 20. Mean, standard deviation has been calculated for age, duration of pain, BMI, and VAS. Frequency and percentage have been calculated for gender, smoking status, diabetes mellitus and MC in the vertebral endplate. Effect



Fig. 2: Modic change at L3



Fig. 3: Modic change at L2



Fig. 1: Modic change at S1

modifier has been controlled through the stratification of age, gender, duration of LBP, BMI, VAS, diabetes mellitus, and smoking status to see the effect of these factors on MC. Post-stratification chi-square test was used, and p-value of  $\leq 0.05$  was considered significant.

## RESULTS

There were a total of 194 patients enrolled in the study, where 58.2% were females, and 41.8% were males. The age of the patients ranged from 30 to 70 years old. Age groups of 30-40 years and 60-70 years were more common, i.e., 27.8% and 29.9% respectively (Fig. 4) with the mean age being 50.36 (SD: 13.22) years (Fig. 5).

Among the enrolled patients, the highest frequency of the patients were farmers (46.4%) followed by 23.7% labors (Fig. 6). The criteria “other” invariably involved the elderly (60-70 years of age), and they were retired office worker, army personnel, or not involved in any activities besides daily chores (Fig. 7).




Fig. 4




Fig. 6




Fig. 5




Fig. 7

In this study, 11.3% of the total sampled patients had a previous history of diabetes mellitus (Table 2), and 42.8% had a habit of smoking (Table 2).

Only those patients with low back pain of more than three months duration were considered in this study. Among them, the most extended period of low back pain was 19 months, with mean duration being 9.64 (SD 2.76) months.

Mean BMI of the patients was 22.50 (SD 2.7) kg/m<sup>2</sup>. Of all the patients, 70.6% of the patients had a healthy weight, and 20.6% of them were overweighted (Table 3).

VAS ranged from 4 to 9 with mean being 6.46 (SD 1.45). Among all the patients, the majority of patients (67.5%) had a moderate degree of pain (Table 4).

Table 2: Demographical data of diabetes and smoking.

Categories		Frequency	Percent
Diabetes Mellitus	No	172	88.7
	yes	22	11.3
Smoking	no	111	57.2
	yes	83	42.8

Table 3: Distribution of patients in different body mass index categories.

BMI categories	Frequency	Percent
Underweight (<18.5)	17	8.8
Healthy weight (between 18.5 and 25)	137	70.6
Over weight (between 25 to 30)	40	20.6

Table 4: Presentation of patients with different severity of pain.

Visual Analogue Scale Categories (VAS)	Frequency	Percent
Moderate pain (VAS 4-7)	131	67.5
Severe pain (VAS >7)	63	32.5
Total	194	100.0

### Prevalence and Association of Factors Affecting Modic Changes

The overall prevalence of MC was 50.6% (Table 5) with MC I being 9.3%, MC II being 35.6% and MC III being 5.7% (Table 6).

**Table 5:** Prevalance of modic changes

Modic changes	Frequency	Percent
No	96	49.5
Yes	98	50.5

**Table 6:** Prevalance of different categories of modic changes.

		Frequency	Percent
	No Modic changes	96	49.5
Modic Changes	Modic I	18	9.3
	Modic II	69	35.6
	Modic III	11	5.7
	Total	194	100.0

**Table 7:** Association of different factors with modic changes.

Categories		Modic changes		P Value
		no change	Modic change	
Gender	Female	60	53	0.248
	Male	36	45	
Age Categories	30-40	40	14	0.000*
	41-50	21	25	
	51-60	12	24	
	61-70	23	35	
Occupation	Labor	24	22	0.123
	Farmer	48	42	
	Office worker	7	17	
	House wife	6	2	
	other	11	15	
VAS	moderate pain	72	59	0.032*
	severe pain	24	39	
Duration of Pain	3 to 6	9	11	0.06
	7 to 10	56	52	
	11 to 14	29	23	
	15 to 18	2	11	
	more than 18	0	1	
DM	no	86	86	0.822
	yes	10	12	
Smoking	no	80	31	0.000*
	yes	16	67	
BMI categories	underweight	12	5	0.001*
	healthy weight	74	63	
	over weight	10	30	

\*means P value <0.05.

MC had no significant gender preponderance (Table 7); however, was significantly associated with age. With advancing age the prevalence of MC was increasing, until the highest prevalence (66.66%) at 61-70 years of age.

Prevalence of MC was highest in patients who have a sedentary life style like office worker (17/24, 70.83%), and retired personnel (15/26, 57.26%); compared to other active groups like farmer and labor. However, the relation was not statistically significant (Table 7).

Prevalence of MC was common if the patient presented with severe pain (39/63, 57.14%) and the value was statistically significant. However, there was no significant association with the duration of pain (Table 7).

DM doesn't have a significant association with MC. But, the patient having a habit of smoking possessed higher prevalence of MC (67/83, 80.72%) with the high level of significance (Table 7).

The higher the BMI category, the higher was the prevalence of MC. The overweighted patients had 75% prevalence of MC which was statistically significant (Table 7).

## DISCUSSION

### Prevalence of Modic Changes

The overall prevalence of MC ranged from 16- 25% in a different study population.<sup>2, 11-15</sup> However in patients with chronic LBP the prevalence was as high as 62%.<sup>16, 17</sup> In this study, 98 out of 194 patients with chronic LBP, who had a history of back pain for more than 3 months, had MC which accounted for 50.5%.

### Modic Changes and Gender

Females seem to have a higher prevalence of MC in most of the studies, which ranged between 52.5 to 70%, but statistic doesn't have a significant association in most of them.<sup>17-22</sup> Similarly, our study also showed a higher prevalence of MC in female though it was statistically not significant.

### Modic Changes and Age

Various studies have included the age from, 18 to 80 years in the study population. And, some of those studies has shown significant association of MC with age.<sup>2, 15-17, 19</sup> In this study, there was a similar finding of increasing prevalence of MC with increasing age groups, and the values were statistically significant.

### Modic Changes and Occupation

Nepal is an agricultural country, 80% of the population survives as a farmer.<sup>23</sup> This evidence was reflected in this study with the maximum number of patients being farmer (90, 46.39%), and was followed by labor (35, 19.4%). Kjaer et al. in their study also showed a relationship between heavy physical work and Modic changes;<sup>24</sup> however, the prevalence of MC was more common in patients who lived a sedentary lifestyle like office worker (70.83%) and retired personnel (57.26%), but the association was not statistically significant.



### Modic Changes and VAS, duration of Pain

This study enrolled the patients with moderate (VAS 4–7) and severe (VAS >7) pain. The prevalence of MC seems to be significantly associated with severity of pain at presentation, 61.9% of MC was noted in the patient who presented with severe pain; however, the prevalence of MC was not associated with the duration of pain.

### Modic Changes and Diabetes, Smoking

Kjaer et al. in their study also showed the positive association of smoking with Modic changes.<sup>24</sup> Similarly, in our study, there was a significant association of smoking with MC in vertebral endplates. The prevalence was 80.7% in patients who had the habit of smoking compared to 27.9% in the non-smoker.

### Modic Changes and BMI

In previous studies, overweight in combination with laborious physical activity was significantly associated with the prevalence of Modic changes.<sup>25</sup> Similar findings were also note in the study of Karchevsky et al. where it showed there was the significant association between prevalence of MC with increasing weight but not with BMI. In this study, the prevalence of MC seems to be increasing from 29.4% among underweighted patients to 75% among overweighted patients with a high level of statistical significance (Table 7)

### CONCLUSION

Modic change occurring at the vertebral endplate is a common phenomenon with the prevalence of 50.5% in patients presented with chronic low back pain, with type II MC having the highest prevalence of all the subtypes. These changes have a significant association with the age, habit of smoking, the severity of pain at presentation, and BMI.

### REFERENCES

- de Roos A, Kressel H, Spritzer C. MR imaging of marrow changes adjacent to end plates in degenerative lumbar disk disease. *AJR Am J Roentgenol.* 1987;149:531–534.
- Modic MT, Steinberg PM, Ross JS. Degenerative disk disease: assessment of changes in vertebral body marrow with MR imaging. *Radiology.* 1988;166:193–199.
- Modic MT, Masaryk TJ, Ross JS. Imaging of degenerative disk disease. *Radiology* 1988;168:177–186.
- Hutton MJ, Bayer JH, Powell JM. Modic Vertebral Body Changes: The Natural History as Assessed by Consecutive Magnetic Resonance Imaging. *Spine.* 2011;36(26):2304–2307.
- Albert HB, Lambert P, Rollason J, et al. Does nuclear tissue infected with bacteria following disc herniations lead to Modic changes in the adjacent vertebrae? *Eur Spine J.* 2013;22(4):690–696.
- Jensen RK, Leboeuf-Yde C, Wedderkopp N, et al. Is the development of Modic changes associated with clinical symptoms? A 14-month cohort study with MRI. *Eur Spine J.* 2012;21(11):2271–2279.
- Sørli A, Moholdt V, Kvistad KA, et al. Modic type I changes and recovery of back pain after lumbar microdiscectomy. *Eur Spine J.* 2012;21(11):2252–2258.
- Kerttula L, Luoma K, Vehmas T, et al. Modic type I change may predict rapid progressive, deforming disc degeneration: a prospective 1-year follow-up study. *Eur Spine J.* 2012;21(6):1135–1142.
- Keller A, Boyle E, Skog TA, et al. Are Modic changes prognostic for recovery in a cohort of patients with non-specific low back pain? *Eur Spine J.* 2012;21(3):418–424.
- Jensen RK, Kjaer P, Korsholm L, et al. The Association Between Vertebral Endplate Signal (Modic) Changes And Low Back Pain In The General Population. *J Bone Joint Surg Br.* 2012;94-B.
- Kuisma M, Karppinen J, Niinimäki J, et al. Modic changes in endplates of lumbar vertebral bodies: prevalence and association with low back and sciatic pain among middle-aged male workers. *Spine.* 2007;32:1116–1122.
- Mitra D, Cassar-Pullicino VN, McCall IW. Longitudinal study of vertebral type-1 end-plate changes on MR of the lumbar spine. *Eur Radiol.* 2004;14(9):1574–1581.
- Toyone T, Takahashi K, Kitahara H, et al. Vertebral bone-marrow changes in degenerative lumbar disc disease. An MRI study of 74 patients with low back pain. *J Bone Joint Surg Br.* 1994;76(5):757–164.
- Weishaupt D, Zanetti M, Hodler J, et al. Painful Lumbar Disk Derangement: Relevance of Endplate Abnormalities at MR Imaging. *Radiology.* 2001;218(2):420–427.
- Albert HB, Manniche C. Modic changes following lumbar disc herniation. *Eur Spine J.* 2007;16(7):977–982.
- Kleinstuck F, Dvorak J, Mannion AF. Are “structural abnormalities” on magnetic resonance imaging a contraindication to the successful conservative treatment of chronic nonspecific low back pain? *Spine.* 2006;31:2250–2257.
- Karchevsky M, Schweitzer ME, Carrino JA, et al. Reactive endplate marrow changes: a systematic morphologic and epidemiologic evaluation. *Skeletal Radiol.* 2005;34(3):125–129.
- Chung CB, VandeBerg BC, Tavernier T, et al. End plate marrow changes in the asymptomatic lumbosacral spine: frequency, distribution and correlation with age and degenerative changes. *Skeletal Radiol.* 2004;33(7):399–404.
- Girard CJ, Schweitzer ME, Morrison WB, et al. Thoracic spine disc-related abnormalities: longitudinal MR imaging assessment. *Skeletal Radiol.* 2004;33(4):216–222.
- Kjaer P, Leboeuf-Yde C, Sorensen JS, et al. An epidemiologic study of MRI and low back pain in 13-year-old children. *Spine.* 2005;30(7):798–806.
- Kjaer P, Leboeuf-Yde C, Korsholm L, et al. Magnetic resonance imaging and low back pain in adults: a diagnostic imaging study of 40-year-old men and women. *Spine.* 2005;30(10):1173–1180.
- Peterson CK, Humphreys BK, Pringle TC. Prevalence of modic degenerative marrow changes in the cervical spine. *J Manipulative Physiol Ther.* 2007;30(1):5–10.
- About Organic Agriculture in Nepal. *SADP Nepal, Sustainable Agriculture Development Program Nepal* 2010.
- Kjaer P, Korsholm L, Bendix T, et al. Modic changes and their associations with clinical findings. *Eur Spine J.* 2006;15(9):1312–1319.
- Leboeuf-Yde C, Kjaer P, Bendix T, et al. Self-reported hard physical work combined with heavy smoking or overweight may result in so-called Modic changes. *BMC Musculoskelet Disord.* 2008;9:5.