

CASE REPORT

Closed Pan-talar Dislocation with Posterior Talar Process Fracture

¹M Pramod Kumar, ²KM Gopinath, ³BN Roshan Kumar, ⁴GA Gautham Balaji

ABSTRACT

Total dislocation of the talus has been reported as 0.06% of all dislocations and 2% of all talar fractures. It usually occurs from considerable violence. Total dislocation of the talus is frequently an open injury, or the skin may be tented over the dislocated talus leading to skin slough. Closed total dislocation of talus with posterior process injury is rare. The functional prognosis is poor due to osteonecrosis of the talus which develops in the majority of cases. Review of literature reported successful closed reduction in cases of closed pan-talar dislocation under anesthesia. However, there were few cases where there was difficulty in closed reduction. Operative technique has also been described in different reports of similar cases.

Case report: We present a case of pan-talar dislocation of the left talus in a 25-year-old road accident victim, with posterior talar process fracture. Reduction of dislocation was attempted in emergency department by external manipulation. Reduction process failed, and hence planned for reduction under anesthesia. It required an open reduction after an unsuccessful closed reduction attempt. The talus after reduction was found to be unstable hence, stabilized with trans-calcaneotalar Steinmann pin. At 1-year follow-up, the right ankle was pain free and stable. Motion was satisfactory. The talus after a follow-up of 1 year did not show any signs of subluxation or avascular necrosis.

Conclusion: The main obstacle to closed reduction appeared to be talus had button holed through dorsal fascia. The talus after reduction was found to be unstable hence stabilized with trans-calcaneotalar Steinmann pin. The management of the associated fracture will depend on many factors, particularly displacement of the fracture fragments.

Keywords: Closed, Dislocation, Operative treatment, Talus.

How to cite this article: Kumar MP, Gopinath KM, Kumar BNR, Balaji GAG. Closed Pan-talar Dislocation with Posterior Talar Process Fracture. *J Med Sci* 2015;1(2):32-35.

Source of support: Nil

Conflict of interest: None

¹Assistant Professor, ²Associate Professor and Unit Head
³Professor and Head, ⁴Postgraduate

¹⁻⁴Department of Orthopedics, RajaRajeswari Medical College and Hospital, Bengaluru, Karnataka, India

Corresponding Author: GA Gautham Balaji, Postgraduate Department of Orthopedics, RajaRajeswari Medical College and Hospital, Bengaluru, Karnataka, India, Phone: 9738220940 e-mail: drgauthambalaji@yahoo.co.in

INTRODUCTION

A closed total dislocation of the talus from all its surrounding joints (talonavicular, tibiotalar, subtalar) not accompanied by a fracture (talus, navicular, calcaneus, malleoli) is an extremely rare injury caused by a high-energy trauma. Its exact incidence is unknown. Closed total talus dislocations are rare.¹ No standard treatment protocol exists regarding treatment of these rare traumatic injuries.² Avascular necrosis is a well-known, severe complication postfracture or subluxation of the talus. Type and localization of injury often permit conclusions regarding the probability of bone necrosis.³

Reviews of case reports open total talar dislocations show subsequent development of infection and avascular necrosis of the talar dome, and poor clinical and functional outcomes associated with this injury. Open total talar dislocation is a relatively rare but debilitating injury. Techniques for managing such injuries include reimplantation of an extruded talus, open reduction internal fixation and fusion (immediate or delayed).^{4,5}

Literature cites infection and avascular necrosis as the main complications associated with a talar extrusion. Good open fracture protocol can reduce the risk of infection. Reduction of the extruded talus is preferable to preserve function and maintain normal hindfoot anatomy. Talcotomy should be reserved as a salvage procedure.⁶

Avascular necrosis is dependant on the amount of blood supply that remains to a bone which is notorious for its problematic vascularity.⁷ We present our method of treatment in this case with closed total talus dislocation without fracture.

CASE REPORT

A 25-year-old male, with alleged history of fall from a two wheeler, presented in our emergency with pain, swelling and deformity of his left ankle.

EXAMINATION

- Swelling over the left foot and ankle and varus deformity of the forefoot (Fig. 1)
- On palpation, bony swelling was present below the lateral malleolus with hollowness below the medial malleolus



- The dorsalis pedis and posterior tibial pulsations were palpated as normal
- Movement of ankle was painful and tender
- No external injury
- No associated injury
- No hypoesthesia over the foot or ankle
- There was no sign of hyperlaxity of other joints.

INVESTIGATIONS

Plain Radiography

Talus was found completely dislocated from the tibio-talar, talocalcaneal and talonavicular joints. The talus lay horizontally just below the lateral malleolus (Figs 2A and B).

Computerized Tomography

Showed pan-talar dislocation with fracture of the posterior process of the talus (Fig. 3).



Fig. 1: Swelling over the left foot and ankle, and varus deformity of the forefoot

Emergency Room

Gentle manipulation was tried twice in the emergency department with the foot in plantar flexion, pronation of foot with knee flexion. Reduction process failed, and hence planned for reduction under anesthesia (Figs 4A and B).

Operative Procedure in the Operation Theater


- Under general anesthesia, reduction was done which also failed.
- Open reduction was done through medial approach. Medial malleolus osteotomy done to approach the talus.
- Open reduction was successful and revealed that the talus had 'button-holed' through the dorsal fascia preventing closed reduction.
- Joint was thoroughly washed.
- Ankle joint was found to be unstable plantigrade, hence a 3.5 mm Steinmann pin passed retrograde from calcaneum to the tibia under fluoroscopy.



Figs 2A and B: Anteroposterior and lateral radiographs of total lateral talar dislocation



Fig. 3: Computerized tomography (CT) showed pan-talar dislocation with fracture of the posterior process of the talus



Figs 4A and B: Anteroposterior and lateral views after closed reduction




Fig. 5: Postoperative lateral X-rays showing the reduced ankle with Steinmann pin and medial malleolus with 4 mm cannulated cancellous screws




Fig. 6: Anteroposterior and lateral views of ankle after the Steinmann pin was removed

- Medial malleolus was fixed with two cannulated cancellous screws. Comminuted posterior process was removed (Fig. 5).
- Postoperatively, immobilization was done with a below knee slab.

Postoperative

- After removing the sutures, a nonweight-bearing below knee cast was applied for 6 weeks.

- After 6 weeks, the Steinmann pin was removed (Fig. 6).
- Left ankle was kept in nonweight-bearing with gradual mobilization for another 6 weeks.

Follow-up

- Patient was reviewed after 6 months and was found to have pain free 10° dorsiflexion and 40° plantar flexion.
- Radiographic examination showed no sign of avascular necrosis of the talus (Figs 7A and B).

DISCUSSION

Total dislocation of the talus has been reported as 0.06% of all dislocations and 2% of all talar fractures. From previous studies, incidence of open pan-talar dislocation was between 10 and 40%. In closed dislocations, significant injury of the soft tissue is present, especially of the skin tented over the head of the talus.⁸

Zimmer and Johnson summarized eight major series plus their experience and reported 79.5% medial, 17% lateral, 2.5% posterior and 1% anterior dislocations. Associated fractures were more frequently seen in lateral dislocations. They reported 25% open dislocations and 4 to 5% avascular necrosis.⁹



Figs 7A and B: (A) 10° dorsiflexion and (B) 40° plantar flexion at 6 months

DeLee and Curtis reported the results of 17 patients involving 70% medial, 23% lateral and 5% anterior dislocations. Open dislocations were reported in 17% and associated fractures involving talocalcaneal and talonavicular joints in 47%.¹⁰

The difficulty in closed reduction that we faced in the present case was reported in many previous studies. In our case, closed reduction was not achieved as there the talus had 'button-holed' through the dorsal fascia preventing reduction. The operative technique discussed in this case was partly described in different reports of similar cases.

CONCLUSION

Total dislocation of talus with posterior process injury is a rare injury, usually an open injury. But, we report a case of closed total dislocation of talus with posterior process fracture, which required an open reduction after an unsuccessful closed reduction attempt. The main obstacle to closed reduction appeared to be talus which had gone through a button hole in the dorsal fascia. The talus after reduction was found to be unstable, hence, stabilized with Steinmann pin. The management of the associated fracture will depend on many factors, particularly displacement of the fracture fragments.

REFERENCES

1. El Ibrahim A, Shimi M, Elidrissi M, Daodi A, Elmrini A. A case of closed total dislocation of talus and literature review. *Am J Emerg Med* 2011;29(4):475. e1-e3.
2. Heylen S, De Baets T, Verstraete P. Closed total talus dislocation: a case report. *Acta Orthop Belg* 2011;77(6):838-842.
3. Schiffer G, Jubel A, Elsner A, Andermahr J. Complete talar dislocation without late osteonecrosis: clinical case and anatomic study. *J Foot Ankle Surg* 2007;46(2):120-123.
4. Ely EE, Konstantakos EK, Laughlin RT, Johnson RM, Binski JC. Total dislocation of the talus and the navicular: a case report. *J Orthop Trauma* 2009;23(7):546-549.
5. Burston JL, Isenegger P, Zellweger R. Open total talus dislocation: clinical and functional outcomes: a case series. *J Trauma* 2011;68(6):1453-1458.
6. Fleming J, Hurley KK. Total talar extrusion: a case report. *J Foot Ankle Surg* 2009;48(6):690. e19-e23.
7. Xarchas KC, Psillakis IG, Kazakos KJ, Pelekas S, Ververidis AN, Verettas DA. Total dislocation of the talus without a fracture open or closed treatment? Report of two cases and review of the literature. *Open Orthop J* 2009;3:52-55.
8. Weston JT, Liu X, Wandtke ME, Liu J, Ebraheim NE. A systematic review of total dislocation of the talus. *Orthop Surg* 2015;7:97-101.
9. Fotiadis E, Lyrtzis C, Svarnas T, Koimtzis M, Akritopoulou K, Chalidis B. Closed subtalar dislocation with non-displaced fractures of talus and navicular: a case report and review of the literature. *Cases J* 2009;2:8793.
10. DeLee JC, Curtis R. Subtalar dislocation of the foot. *J Bone and Joint Surg* 1982 Mar 1;64(3):433-437.