

Postoperative Nausea and Vomiting in Day Care Patients: A Comparative Randomized Controlled Trial of Total Intravenous Anesthesia with Propofol, Air, and Oxygen vs Inhalation Anesthesia with Isoflurane and Nitrous Oxide

¹A Sandhya, ²R Mamatha, ³Antara Banerjee, ⁴H Sahajananda

ABSTRACT

Introduction: We compared the incidence of postoperative nausea and vomiting (PONV) after total intravenous anesthesia (TIVA) using propofol–air to inhalational anesthesia with isoflurane–nitrous oxide in day care patients at a tertiary care academic institution.

Materials and methods: We randomized 60 patients and assigned to either group I (inhalational anesthesia with isoflurane–nitrous oxide) or group II (TIVA with propofol–air). Incidence of PONV, use of anti-emetics, and duration of stay in the recovery were recorded for 72 hours by blinded observers.

Results: Total intravenous anesthesia reduced the PONV up to 72 hours by 27% among our patients (from 37 to 10%, $p < 0.001$). This effect was seen more in the early postoperative period. Overall, 13.3% of patients in the group I received antiemetic compared to 40% in group II. In our study, patients without PONV were discharged from the recovery room 15 minutes earlier after TIVA than after isoflurane and N₂O anesthesia.

Conclusion: Total intravenous anesthesia with propofol and air resulted in a reduction of PONV compared with isoflurane–nitrous oxide anesthesia. Overall, patients in group I required less rescue antiemetic, compared to group II. Total intravenous anesthesia resulted in shorter stay in the postoperative anesthetic care unit compared to isoflurane–N₂O group.

Keywords: Inhalational anesthesia, Postoperative nausea and vomiting, Total intravenous anesthesia.

How to cite this article: Sandhya A, Mamatha R, Banerjee A, Sahajananda H. Postoperative Nausea and Vomiting in Day Care Patients: A Comparative Randomized Controlled Trial of Total Intravenous Anesthesia with Propofol, Air, and Oxygen vs Inhalation Anesthesia with Isoflurane and Nitrous Oxide. *J Med Sci* 2015;1(4):63-68.

Source of support: Nil

Conflict of interest: None

¹⁻³Postgraduate Student, ⁴Professor and Head

¹⁻⁴Department of Anesthesiology, RajaRajeswari Medical College and Hospital, Bengaluru, Karnataka, India

Corresponding Author: H Sahajananda, Professor and Head, Department of Anesthesiology, RajaRajeswari Medical College and Hospital, Bengaluru, Karnataka, India, Phone: +919448085401, e-mail: sahaj_anand@hotmail.com

INTRODUCTION

The use of total intravenous anesthesia (TIVA) using propofol alone or inhalational anesthesia using isoflurane with air and oxygen in day care surgeries is always a matter of concern.¹ Studies have shown that TIVA with propofol reduces the incidence of postoperative nausea and vomiting (PONV) and results in shorter recovery time.²⁻⁶ Most of those studies were smaller, did not have follow-up more than 6 hours postoperatively, and many of the studies were sponsored by pharmaceutical companies. The results were difficult to substantiate as a result of multiple definitions of PONV.^{7,8}

Hence we compared the incidence of PONV up to 72 hours postoperatively in outpatients who were randomized to receive either inhalational anesthesia with isoflurane–N₂O or TIVA with propofol–air. The primary aim and hypothesis was that propofol TIVA would probably reduce the incidence of PONV compared with an inhalation anesthetic technique using isoflurane and N₂O.

MATERIALS AND METHODS

We conducted a randomized controlled trial of TIVA with propofol vs inhalational anesthesia with isoflurane–air and oxygen. The study was conducted at RajaRajeswari Medical College and Hospital, Bengaluru and was approved by the Institutional Medical Ethics Committee. Sixty patients scheduled to undergo elective surgery under general anesthesia were enrolled in the study. Exclusion criteria were emergency, cardiac, or neurosurgery, American Society of Anesthesiologists physical status classification greater than III, age < 18 years or > 80 years, pregnancy, renal or liver disease, use of antiemetic in the 2 weeks before surgery, and body weight more than 120 kg.

PREOPERATIVE EVALUATION

Preoperative evaluation was done 1 day before the surgery. This involved detailed history from the patient, and any medical complaints that the patient had also noted. General physical examination was done along with a detailed evaluation of the cardiovascular, respiratory,

and central nervous system. All routine and relevant lab investigations were carried out.

PREOPERATIVE ORDER

Patients were instructed to stay nil per oral after 10 pm. All of them received Tab. alprazolam 0.25 mg orally on the night before surgery. They were given Tab. ranitidine 150 mg orally on the morning of surgery with sips of water and Inj. midazolam 1 mg intravenously (IV) 1 hour before surgery.

All patients were allocated into two groups using computerized randomization table.

Group I: Total intravenous anesthesia with propofol and air.

Group II: Inhalation with isoflurane, nitrous oxide, and oxygen.

Standard monitors were attached along with bispectral index (BIS) monitoring.

In the study (TIVA) group I, general anesthesia was induced with propofol 2 to 3 mg/kg and maintained on a continuous infusion of 4 to 12 mg/kg/hour, with 60% nitrous oxide and 33% oxygen mixture. The rate of propofol infusion was adjusted according to the need of anesthesia depth by using BIS. Propofol was tapered toward the end of case and stopped at skin closure. Both groups received 0.5 mg/kg atracurium for intubation and further maintained on one-third of the original dose, as top-up doses. Intraoperatively, analgesia in both the groups was provided with 2 µg of fentanyl at induction, along with multimodal analgesia using paracetamol 1 g IV, 75 mg of Inj. diclofenac, and infiltration analgesia before skin incision.

In this study, the symptoms of PONV, such as nausea, retching, and vomiting, were observed and the severity of PONV was rated at timely intervals. Use of antiemetics, pain scores [visual analog scale (VAS) score], patient satisfaction scores, and time of discharge from the recovery area were also noted.

Monitoring

Intraoperative heart rate (HR), blood pressure (BP), EtCO₂, and electrocardiogram (ECG) were monitored continuously. The duration of surgery was noted.

Postoperatively, patients were observed for 24, 48, and 72 hours for nausea, retching, and vomiting.

Pain was assessed using VAS score; in case VAS score exceeded 4/5, patients were given rescue analgesia as Inj. diclofenac 75 mg IM.

Rescue antiemetic comprised 0.15 mg/kg metoclopramide IV and was given if patients were having retching. Rescue antiemetic doses if given were also noted. Complications if any were also noted.

Postoperative Assessment: Measurements

Routine monitoring was initiated on arrival in the postoperative anesthetic unit (PACU) or day care unit. Every 15 minutes, anesthesiologist recorded PONV, use of analgesic, and antiemetic medication. Nausea, retching, and vomiting were scored separately.

In the postoperative period, the patients were monitored for the 1st 24 hours:

They were monitored for:

- Episodes of nausea: Defined as a subjective unpleasant feeling of having to vomit. This was scored 0 to 3.
0 = no nausea
1 = mild nausea
2 = moderate nausea
3 = severe nausea
- Episode of retching was defined as vomiting movement without actual vomiting, which was scored as absent or present.
- They were also observed for episodes of vomiting, that is expulsion of stomach contents, and whether they had early or late PONV.
- A cumulative PONV score was calculated taking into account nausea, vomiting, and retching scores.
- The patients were also assessed for pain using the VAS for the 1st 24 hours.
- Approximately 24 hours after surgery, an anesthesiologist visited patients in the ward or telephoned outpatients at home to record occurrence of PONV, use of antiemetics or analgesics, and the presence of possible postoperative complications and side effects subsequent to discharge from the PACU. The time of discharge and duration of stay in the recovery area were also noted. Patients were asked to quantify discomfort caused by nausea, retching, and vomiting and to rate their anesthetic experience (in the scale of 0–10). Similar information was collected 72 hours postoperatively with reference to the previous 48 hours.

STATISTICAL ANALYSIS

Data were collected and entered in MS Excel and analyzed using Statistical Package for the Social Sciences (SPSS) version 2.0. Descriptive statistics includes frequencies, percentages, and mean with standard deviation. Student's t-test will be used to test the significant difference between the two groups with 95% confidence interval.

Sample size is calculated to be 66; 33 in each group by the formula

$$n = 4 PQ/L^2$$

where P = prevalence of PONV in gynecological patients undergoing laparoscopic surgeries, i.e., 75%; Q = 100–P; L = allowable error. (20% of P).

RESULTS

Six patients were excluded from the study as they could not be contacted for the follow-up after 24 hours.

Baseline patient characteristics were similar across allocation groups (Table 1). The average age of patients was 60.10 and 59.20 years respectively. The mean BMI in group I was 24.85 ± 4.75 as against 24.66 ± 3.42 in group II with a p value of 0.856, which was statistically insignificant.

Types of Surgery

In all the groups duration of anesthesia and surgery were also similar and not significant (Graphs 1 and 2). The mean duration of surgery was 87.50 ± 31.07 in group I as against 87.44 ± 27.47 in group II. Average duration of

surgery for both groups was 60 to 90 minutes. This was found to be statistically insignificant.

In group I out of a total of 30, 16 patients underwent superficial surgeries, 9 underwent Diagnostic Hysteroscopic Laparoscopy, 3 underwent cystoscopy, 1 underwent salpingectomy and 1 underwent cystectomy. While in group II out of the total 30, 15 patients underwent superficial surgeries, 13 underwent Diagnostic Hysteroscopic Laparoscopy, 1 underwent cystoscopy, 1 underwent salpingectomy. The p-value for the type of surgery done was not found to be significant (Graph 3).


The mean dose of fentanyl in group I was 101.33 ± 17.95 as compared to 97.67 ± 14.06 in group II, this was found to have a p value of 0.382, which was not statistically significant (Graph 4).

Hence the two groups were well randomized and statistically comparable in terms of age, sex, weight, duration of surgery, anesthesia, types of surgery, and opioid use in our study.


Endotracheal tubes and muscle relaxants were used for all patients. All patients received opioids intraoperatively.

Table 1: Demographic data – mean (SD)


	Group I	Group II	p-value
Age (years)	60.10 ± 13.60	59.20 ± 9.64	0.769
Height (cm)	155.03 ± 5.49	154.73 ± 5.72	0.837
BMI (kg/m ²)	24.85 ± 4.75	24.66 ± 3.42	0.856




Graph 1: Duration of surgery (minutes)



Graph 2: Duration of anesthesia (minutes)



Graph 3: Types of surgery



Graph 4: Fentanyl dose (µg)

The average use of propofol for anesthesia maintenance was 10.0 mg/kg/hour. Isoflurane use was 12.5 mL/hour. The time from discontinuation of anesthesia until response to a verbal command in outpatients after TIVA or isoflurane was 9 minutes.

Postoperative Nausea and Vomiting

Nausea

In group I, 10% of patients had nausea compared to group II who had 36.7% in the 1st 24 hours. It was statistically significant (p=0.029). In the next 48 hours and at 72 hours, the incidence of nausea was not significant (Table 2).

Retching

Incidence of retching between the two groups was not significant over 24 hours (Table 3).

Vomiting

Incidence of vomiting was more in group I compared to group II at 24 hours and statistically it was significant (Table 4); whereas at 48 and 72 hours incidence, it was not significant.

Table 2: Incidence of nausea in numbers (percentages)

Nausea	24 hours	48 hours	72 hours	% Change
Group I (n=30)				
0	27 (90%)	28 (93.3%)	30 (100%)	10.0
1	1 (3.3%)	0 (0%)	0 (0%)	0.0
2	0 (0%)	2 (6.7%)	0 (0%)	-3.3
3	2 (6.7%)	0 (0%)	0 (0%)	-6.7
Group II (n=30)				
0	19 (63.3%)	29 (96.7%)	30 (100%)	36.7
1	1 (3.3%)	0 (0%)	0 (0%)	-23.3
2	6 (20%)	0 (0%)	0 (0%)	-13.3
3	4 (13.3%)	1 (3.3%)	0 (0%)	0.0
p-value	0.029*	0.492	1.000	-

* Moderately Significant

Table 4: Incidence of vomiting numbers (percentages)

Vomiting	24 hours	48 hours	72 hours
Group II (n=30)			
No	25 (83.33%)	28 (93.33)	29 (96.7%)
Yes	5 (16.66%)	2 (6.67%)	1 (3.3%)
Group I (n=30)			
No	28 (93.33%)	29 (96.7)	29 (96.7%)
Yes	2 (6.67%)	1 (3.3%)	1 (3.3%)
p-value	0.001	0.453	1.000

PONV Score

The overall incidence of vomiting between the two groups was significant at 24 hours (p=0.001). It was not significant at 48 and 72 hours (Table 5).

Rescue Antiemetic

Overall, four patients in group I required rescue antiemetic, while 12 in group II needed rescue antiemetic. Requirement of rescue antiemetic between the two groups was not significant statistically (Table 6).

Pain

Following surgery, the pain scores between the two groups were comparable, and not statistically significant.

DISCUSSION

Postoperative nausea and vomiting are common and distressing to patients, with an incidence of as high as 80% in high-risk patients. According to the 2014 consensus guidelines for the management of PONV by the American Society of Anesthesiologist, the first step in management is identification of high-risk individuals. Of the various scoring systems, the Apfel risk score is one of

Table 3: Incidence of retching numbers (percentages)

Retching	0-24 hours	48 hours	72 hours	% Change
Group I (n=30)				
No	28 (93.3%)	27 (90%)	28 (93.3%)	10.0
Yes	2 (6.7%)	3 (10%)	2 (6.7%)	-10.0
Group II (n=30)				
No	23 (76.7%)	30 (100%)	26 (86.7%)	0.0
Yes	7 (23.3%)	0 (0%)	4 (13.3%)	0.0
p-value	0.145	0.237	0.671	-

Table 5: Incidence of PONV in numbers (percentages)

PONV score	24 hours	48 hours	72 hours	% Change
Group I (n=30)				
0	27 (90%)	28 (93.3%)	30 (100%)	6.7
1	0 (0%)	0 (0%)	0 (0%)	0.0
2	0 (0%)	0 (0%)	0 (0%)	0.0
3	0 (0%)	1 (3.3%)	0 (0%)	-3.3
4	2 (6.7%)	0 (0%)	0 (0%)	0.0
5	1 (3.3%)	1 (3.3%)	0 (0%)	-3.3
Group II (n=30)				
0	19 (63.3%)	23 (76.7%)	28 (93.3%)	20.0
1	7 (23.3%)	0 (0%)	0 (0%)	-13.3
2	0 (0%)	3 (10%)	0 (0%)	-13.3
3	0 (0%)	0 (0%)	0 (0%)	0.0
4	0 (0%)	1 (3.3%)	1 (3.3%)	3.3
5	4 (13.3%)	3 (10%)	1 (3.3%)	3.3
p-value	0.001*	0.150	0.492	-

*Highly significant

Table 6: Use of antiemetics in numbers and percentages

Rescue antiemetic	Group I (n=30)		Group II (n=30)		p-value
	No	%	No	%	
24 hours	1	3.3	2	6.7	1.000
48 hours	1	3.3	4	13.3	1.000
72 hours	1	3.3	4	13.3	1.000
	4	13.3	12	40	

the commonly used ones. It is based on four predictors, such as female sex, nonsmokers, patients with a previous history of PONV, and who are on opioid drugs. Patients with three or more risk factors are considered at high risk for PONV. Our study included a moderate- to high-risk population, i.e., females in the age group 25 to 40 years who are nonsmokers and not on any opioids.

Strategies to reduce the baseline risk are by using regional anesthesia over general, propofol induction, avoiding nitrous oxide and volatile anesthetics, minimizing the use of opioids perioperatively and use of good hydration. Patients in our study were all induced with propofol and a single dose of fentanyl at induction. General anesthesia, volatile agents like isoflurane and use of nitrous oxide were factors which could not be avoided.

We compared and studied the incidence of PONV up to 72 hours postoperatively in day care patients at a tertiary care academic institution who were randomized to receive either inhalational anesthesia with isoflurane-N₂O or TIVA with propofol-air.

Our study and control groups were both comparable demographically. Duration of anesthesia (Graphs 1 to 3) and intraoperative (Graph 4) and post operative analgesia (Table 7) was comparable between the two groups.

In our study, the incidence of PONV was significantly lower after TIVA than after isoflurane. In TIVA group I, absolute risk reduction was between 27%, which depends on the duration of follow-up. Moreover, in patients' view, TIVA was superior to isoflurane with N₂O. The reduction in PONV in our study is in agreement with results from two recent meta-analyses by Tramer et al⁷ and Sneyd et al⁸ comparing propofol with isoflurane inhalational agent. Follow-up period in our study was long enough compared with few other PONV studies. We think that the anesthetic technique in the 1st 24 hours after surgery, probably, has reduced the incidence of PONV. At 48 and 72 hours, the incidence of PONV increased equally in both groups, hence suggests that

Table 7: Mean visual analog score (SD)

Pain	Group I	Group II	p-value
24 hours	3.17±1.05	3.17±0.99	1.000
48 hours	3.63±0.96	4.03±0.89	0.100
72 hours	3.00±0.95	3.17±1.05	0.522

the anesthetic technique influences the PONV in the 1st 24 hours after surgery, whereas PONV resulting from the surgery and analgesics in the postoperative period dominates thereafter.

Various definitions of PONV have been used, e.g., retching, nausea only, nausea and vomiting, or vomiting only. This in turn has led to difficulties in comparing the various studies.^{9,10} In our study, we scored nausea, retching, and vomiting separately.

Antiemetics were administered more often in patients who received isoflurane than TIVA, 40 vs 13.3%. Subjective ratings of patient satisfaction were highest at 24 hours in the TIVA group. Even after 48 and 72 hours, the ratings of patient satisfaction in general were higher for TIVA than for isoflurane.

All our patients received propofol for induction. Propofol used for anesthesia induction has antiemetic property. This should have reduced the incidence of PONV, irrespective of the maintenance regimen. It is not the case in the present study. This finding is in agreement with the results from a meta-analysis by Tramer et al.⁷

They showed that propofol for anesthesia induction followed by a nonpropofol maintenance technique did not result in the reduction of PONV. Many authors have used N₂O during propofol anesthesia. The use of N₂O allows for lower propofol infusion rates^{11,12} in TIVA and reduces the incidence of awareness during anesthesia and intraoperative period.¹³ It is said that N₂O may increase the incidence of PONV.¹⁴ A recent meta-analysis by Divatia et al¹⁵ has also substantiated this view.

In our study protocol, patients are required to remain in the PACU for at least 1 hour. We found that patients without PONV were discharged from the recovery room 15 minutes earlier after TIVA than after isoflurane. This has also been reported by other investigators.^{16,17}

DRAWBACKS OF THE STUDY

This study was conducted in a single academic institution and these results need to be validated. Further, large-scale clinical trials are required.

CONCLUSION

Total intravenous anesthesia with propofol compared to isoflurane with N₂O was associated with a significantly reduced rate of PONV in the 1st 24 hours and reduced antiemetic consumption. Total intravenous anesthesia increases patient comfort and patient ratings of anesthesia, while slightly reducing discharge time.

REFERENCES

1. Visser K, Hassink EA, Bonsel GJ, Moen J, Kalkman CJ. Randomized controlled trial of total intravenous anesthesia

- with propofol *vs* inhalation anesthesia with isoflurane–nitrous oxide: postoperative nausea and vomiting and economic analysis. *Anesthesiology* 2001 Sep;95(3):616-626.
2. Borgate A, Wilder-Smith OH, Saiah M, Rifat K. Subhypnotic doses of propofol possess direct antiemetic properties. *Anesth Analg* 1992 Apr;74(4):539-541.
 3. Kalman SH, Jensen AG, Ekberg K, Eintrei C. Early and late recovery after major abdominal surgery: comparison between propofol anaesthesia with and without nitrous oxide and isoflurane anaesthesia. *Acta Anaesthesiol Scand* 1993 Nov;37(8):730-736.
 4. Raftery S, Sherry E. Total intravenous anaesthesia with propofol and alfentanil protects against postoperative nausea and vomiting. *Can J Anaesth* 1992 Jan;39(1):37-40.
 5. Oddby-Muhrbeck E, Jakobsson J, Andersson L, Askergrén J. Postoperative nausea and vomiting: a comparison between intravenous and inhalation anaesthesia in breast surgery. *Acta Anaesthesiol Scand* 1994 Jan;38(1):52-65.
 6. Klockgether-Radke A, Piorek V, Crozier T, Kettler D. Nausea and vomiting after laparoscopic surgery: a comparison of propofol and thiopentone/halothane anaesthesia. *Eur J Anaesthesiol* 1996 Jan;13(1):3-9.
 7. Tramer M, Moore A, McQuay H. Propofol anaesthesia and postoperative nausea and vomiting: quantitative systematic review of randomized controlled studies. *Br J Anaesth* 1997 Mar;78(3):247-255.
 8. Sneyd JR, Carr A, Byrom WD, Bilski AJ. A meta-analysis of nausea and vomiting following maintenance of anaesthesia with propofol or inhalational agents. *Eur J Anaesthesiol* 1998 Jul;15(4):433-445.
 9. Korttila K. The study of postoperative nausea and vomiting. *Br J Anaesth* 1992;69(7 Suppl 1):20S-23S.
 10. Andrews PL. Physiology of nausea and vomiting. *Br J Anaesth* 1992;69(7 Suppl 1):2S-19S.
 11. Davidson JA, Macleod AD, Howie JC, White M, Kenny GN. Effective concentration 50 for propofol with and without 67% nitrous oxide. *Acta Anaesthesiol Scand* 1993 Jul;37(5):458-464.
 12. Raeder JC. Total intravenous anaesthesia – free from nitrous oxide, free problems? *Acta Anaesthesiol Scand* 1994 Nov;38(8):769-770.
 13. Tramer M, Moore A, McQuay H. Omitting nitrous oxide in general anaesthesia: meta-analysis of intraoperative awareness and postoperative emesis in randomized controlled trials. *Br J Anaesth* 1996 Feb;76(2):186-193.
 14. James MFM. Nitrous oxide: still useful in the year 2000? *Curr Opin Anaesthesiol* 1999 Jul;12(4):461-466.
 15. Divatia JV, Vaidya JS, Badwe RA, Hawaldar RW. Omission of nitrous oxide during anesthesia reduces the incidence of postoperative nausea and vomiting: a meta-analysis. *Anesthesiology* 1996 Nov;85(5):1055-1062.
 16. Boldt J, Muller M, Uphus D, Hempelmann G. Cost aspects in anesthesia: propofol *vs* isoflurane anesthesia [Article in German]. *Anaesthetist* 1996 Aug;45(8):731-736.
 17. Phillips AS, Mirakhur RK, Glen JB, Hunter SC. Total intravenous anaesthesia with propofol or inhalational anaesthesia with isoflurane for major abdominal surgery: recovery characteristics and postoperative oxygenation – an international multicentre study. *Anaesthesia* 1996 Nov;51(11):1055-1059.