

REHAB QUIZ

1. **All are true about Fredrick ataxia except**
 - A) Nystagmus
 - B) Pes cavus
 - C) Scoliosis
 - D) Ophthalmoplegia
2. **Which is not true about unclassified spondyloarthropathy (USPA)**
 - A) Females are common victims
 - B) Onset in around 5th decade
 - C) Mostly HLA B27 positive
 - D) Dactylitis
3. **In fast walking, which one of the following is correct?**
 - A) The trunk is inclined forward
 - B) The period of double support is longer
 - C) The step length is shorter
 - D) The cadence is increased
4. **Which of the following is not characteristic of F wave?**
 - A) Pathway involving only a motor fibre
 - B) Does not disappear when a motor response is elicited
 - C) Variable latency
 - D) Longer latency than the H reflex of the same nerve muscle stimulation.
5. **The rigid dressing applied with immediate post surgical prosthesis benefits the patients mainly by**
 - A) Assisting the patient to regain a better gait pattern.
 - B) Assisting in primary closure of wound
 - C) Preventing the swelling of stump
 - D) Eliminating the need of a surgical drain
6. **Total hip arthroplasty is contraindicated in all of the following conditions except**
 - A) Paretic gluteus medius
 - B) Avascular necrosis
 - C) Septic arthritis
 - D) Neuropathic joint
7. **Port wine facial nevi, convulsion, calcified areas in cortex suggest the diagnoses of**
 - A) Hemangioblastomas
 - B) Von Recklinghausen's disease
 - C) Sturge- Weber disease
 - D) Tuberos sclerosis
8. **The proprioceptive facilitation technique of irradiation requires**
 - A) Passive movement
 - B) Extrinsic stimulus
 - C) Stretching
 - D) Resistance
9. **Christmas tree appearance in fluoroscope is noticed in**
 - A) Transforaminal epidural block
 - B) Caudal epidural block
 - C) Facet joint block
 - D) Intralaminar epidural block.
10. **All are true regarding the doses guideline of botulinium toxin A for children except**
 - A) Total maximum body dose per visit is 400 IU
 - B) Maximum dose per large muscle per visit is 6 unit per kg
 - C) Maximum dose per injection site is 80 unit.
 - D) Reinjection is ≥ 3 months

ANSWERS

December 2013 issue:

1-B; 2-A; 3-D; 4-A; 5-D; 6-B; 7-D; 8-C; 9-B; 10-C